

Online article and related content
current as of July 11, 2009.

Supplementary material	Spanish PDF http://jama.ama-assn.org/cgi/content/full/288/1/128/DC1
Correction	Contact me if this article is corrected.
Citations	Contact me when this article is cited.
Topic collections	Bacterial Infections; JAMA Patient Page; Infectious Diseases Contact me when new articles are published in these topic areas.
Related Articles published in the same issue	Changing Profile of Infective Endocarditis: Results of a 1-Year Survey in France Bruno Hoen et al. JAMA. 2002;288(1):75.

Subscribe
<http://jama.com/subscribe>




Permissions
permissions@ama-assn.org
<http://pubs.ama-assn.org/misc/permissions.dtl>

Email Alerts
<http://jamaarchives.com/alerts>

Reprints/E-prints
reprints@ama-assn.org

Endocarditis

Endocarditis is an infection or inflammation (response to injury of the cells) of the valves of the heart or of the **endocardium**, the inner lining of the heart chambers. Endocarditis is a serious condition that can cause heart failure, stroke, kidney failure, and death. Fortunately, it can be successfully treated in most people. The July 3, 2002, issue of JAMA includes an article about endocarditis.



TYPES OF ENDOCARDITIS

Endocarditis typically occurs when bacteria enter the bloodstream and then settle on the heart valves. Individuals who have preexisting heart valve problems and persons who have had heart valve operations are at increased risk. Some people get endocarditis from bacteria in the mouth that enter the bloodstream during dental procedures. Unsterile injections by drug abusers also can release bacteria into the blood. For individuals with increased risk, taking antibiotics before medical and dental procedures may reduce the chances of getting endocarditis.

- **Bacterial endocarditis** is caused by bacteria that have entered the bloodstream.
- **Fungal endocarditis** usually occurs in individuals who have low resistance to infection, such as those who are taking medications that suppress the immune system.
- **Noninfective endocarditis** is inflammation in the heart that can be caused by advanced stages of cancer and by some disorders of the immune system.

SYMPTOMS OF ENDOCARDITIS

- Fever and chills
- Drenching night sweats
- Fatigue
- Weight loss
- Shortness of breath and persistent cough
- Pain in the joints
- Bleeding under the fingernails, causing red lines
- **Petechiae**, which are tiny purple-red spots under the skin

TREATMENT

Bacterial endocarditis is treated with antibiotics, usually **intravenously** (administered through the veins) and usually for an extended period (up to 6 weeks). In severe cases in which the heart valves become damaged, surgery may be necessary.

FOR MORE INFORMATION

- American Heart Association
800/242-8721
www.americanheart.org
- National Heart, Lung, and Blood Institute
(National Institutes of Health)
301/592-8573
www.nhlbi.nih.gov

INFORM YOURSELF

To find this and previous JAMA Patient Pages, go to the Patient Page Index on JAMA's Web site at www.jama.com. A Patient Page on heart failure was published in the February 6, 2002, issue.

Sources: American College of Cardiology, American Heart Association, National Heart, Lung, and Blood Institute

Lise M. Stevens, MA, Writer

Cassio Lynn, MA, Illustrator

Richard M. Glass, MD, Editor

The JAMA Patient Page is a public service of JAMA. The information and recommendations appearing on this page are appropriate in most instances, but they are not a substitute for medical diagnosis. For specific information concerning your personal medical condition, JAMA suggests that you consult your physician. This page may be reproduced noncommercially by physicians and other health care professionals to share with patients. Any other reproduction is subject to AMA approval. To purchase bulk reprints, call 718/946-7424.

