



Online article and related content  
current as of November 27, 2009.

## Weight Loss With Self-help Compared With a Structured Commercial Program: A Randomized Trial

Stanley Heshka; James W. Anderson; Richard L. Atkinson; et al.

JAMA. 2003;289(14):1792-1798 (doi:10.1001/jama.289.14.1792)

<http://jama.ama-assn.org/cgi/content/full/289/14/1792>

Correction

[Contact me if this article is corrected.](#)

Citations

[This article has been cited 77 times.](#)  
[Contact me when this article is cited.](#)

Topic collections

Public Health; Obesity; Randomized Controlled Trial  
[Contact me when new articles are published in these topic areas.](#)

Subscribe

<http://jama.com/subscribe>

Permissions

[permissions@ama-assn.org](mailto:permissions@ama-assn.org)  
<http://pubs.ama-assn.org/misc/permissions.dtl>

Email Alerts

<http://jamaarchives.com/alerts>

Reprints/E-prints

[reprints@ama-assn.org](mailto:reprints@ama-assn.org)

# Weight Loss With Self-help Compared With a Structured Commercial Program

## A Randomized Trial

Stanley Heshka, PhD

James W. Anderson, MD

Richard L. Atkinson, MD

Frank L. Greenway, MD

James O. Hill, PhD

Stephen D. Phinney, MD, PhD

Ronette L. Kolotkin, PhD

Karen Miller-Kovach, MS, RD

F. Xavier Pi-Sunyer, MD

**A**N INCREASING NUMBER OF MEN and women in the United States are attempting to lose weight.<sup>1,2</sup> Formal diet and exercise programs provide weight loss treatment to approximately 13% of these women and 5% of the men.<sup>3</sup> However, obese participants in diet and exercise programs rarely achieve a body mass index (BMI, calculated as weight in kilograms divided by the height in meters squared) in the normal range (20-25) and because weight lost is often regained, there is a perception that nothing is accomplished by participation in such programs. Nevertheless, numerous studies have documented improvements in indicators of health with modest weight loss,<sup>4-7</sup> which, if maintained over several years, may be clinically significant.

Commercial weight loss programs have been recommended as the second step in a graded, 4-step model for weight management<sup>5,8</sup> but their efficacy has rarely been rigorously evaluated.<sup>9-13</sup> We conducted a 2-year, mul-

**Context** Although commercial weight loss programs provide treatment to millions of clients, their efficacy has not been evaluated in rigorous long-term trials.

**Objective** To compare weight loss and health benefits achieved and maintained through self-help weight loss vs with a structured commercial program.

**Design and Setting** A 2-year, multicenter randomized clinical trial with clinic visits at 12, 26, 52, 78, and 104 weeks conducted at 6 academic research centers in the United States between January 1998 and January 2001.

**Participants** Overweight and obese men (n=65) and women (n=358) (body mass index, 27-40) aged 18 to 65 years.

**Intervention** Random assignment to either a self-help program (n=212) consisting of two 20-minute counseling sessions with a nutritionist and provision of self-help resources or to a commercial weight loss program (n=211) consisting of a food plan, an activity plan, and a cognitive restructuring behavior modification plan, delivered at weekly meetings.

**Main Outcome Measures** Weight change was the primary outcome measure. Secondary outcomes included waist circumference, body mass index, blood pressure, serum lipids, glucose, and insulin levels.

**Results** At 2 years, 150 participants (71%) in the commercial group and 159 (75%) in the self-help group completed the study. In the intent-to-treat analysis, mean (SD) weight loss of participants in the commercial group was greater than in the self-help group at 1 year (-4.3 [6.1] kg vs -1.3 [6.1] kg, respectively;  $P < .001$ ) and at 2 years (-2.9 [6.5] kg vs -0.2 [6.5] kg, respectively;  $P < .001$ ). Waist circumference ( $P = .003$ ) and body mass index ( $P < .001$ ) decreased more in the commercial group. Changes in blood pressure, lipids, glucose, and insulin levels were related to changes in weight in both groups, but between-group differences in biological parameters were mainly nonsignificant by year 2.

**Conclusion** The structured commercial weight loss program provided modest weight loss but more than self-help over a 2-year period.

JAMA. 2003;289:1792-1798

www.jama.com

**Author Affiliations:** New York Obesity Research Center, St Luke's/Roosevelt Hospital and Columbia University, New York, NY (Drs Heshka and Pi-Sunyer); VA Medical Center, Lexington, and University of Kentucky (Dr Anderson); Nutritional Sciences, University of Wisconsin, Madison (Dr Atkinson); Pennington Biomedical Research Center, Louisiana State University, Baton Rouge (Dr Greenway); Health Sciences, University of Colorado, Boulder (Dr Hill); Department of Medicine, University of California at Davis (Dr Phinney); Obesity and Quality of Life Consulting, Durham, NC (Dr Kolotkin); and Weight Watchers International, Woodbury, NY (Ms Miller-Kovach).

**Financial Disclosures:** Dr Anderson has received grants/research support from Abbott, AstraZeneca, Aventis, Bristol-Myers Squibb, GlaxoSmithKline, Health

Management Resources, Johnson & Johnson, Kos, Merck, NutriPharma, Protein Technologies International, Procter & Gamble, Regeneron, Roche, Schering-Plough, Tishcon, Weight Watchers, and Wyeth-Ayerst, has served as a consultant to GlaxoSmithKline, Health Management Resources, Johnson & Johnson, Knoll, NutriPharma, and Protein Technologies International, and has received honoraria from Abbott, Bristol-Myers Squibb, GlaxoSmithKline, NutriPharma, and Protein Technologies International; Dr Greenway is a consultant to Jenny Craig; and Dr Pi-Sunyer serves on the scientific advisory board for Weight Watchers.

**Corresponding Author and Reprints:** Stanley Heshka, PhD, St Luke's/Roosevelt Hospital Center, 1090 Amsterdam Ave, 14C, New York, NY 10025 (e-mail: sh311@columbia.edu).

ticenter randomized trial to evaluate the largest provider of commercial weight loss services in the United States.<sup>14</sup> A report of the 6-month results of this trial has been published previously.<sup>15</sup>

## METHODS

### Study Design

The study was a randomized, parallel-group, 2-year trial conducted at 6 US clinical centers (see author affiliations on the first page) between January 1998 and January 2001 with scheduled visits to the study center at weeks 0, 12, 26, 52, 78, and 104. The number of participants to be enrolled was selected to provide a power of 0.84 to detect a difference of 2.5 kg between treatment and control groups in the amount of weight lost at the end of 2 years based on an estimated SD of 6.5 kg in change scores, with a type I error rate of .05, 2-tailed, if 120 participants remained in each group.

### Study Participants

Participants were recruited from existing clinic records or by advertising a long-term nonmedication weight loss study for moderately overweight persons. All participants provided written informed consent. A medical history was taken and physical examination and electrocardiography were performed at the screening visit. Men and women with a BMI of 27 to 40, aged 18 to 65 years, including persons with health problems for which weight reduction is a medically accepted therapy, were eligible for the study.

Exclusion criteria were fasting glucose higher than 140 mg/dL (7.8 mmol/L), triglycerides higher than 1000 mg/dL (11.3 mmol/L), liver function test results (aspartate aminotransferase, alanine aminotransferase, alkaline phosphatase, lactate dehydrogenase,  $\gamma$ -glutamyltransferase, and bilirubin) more than 2 times the upper normal limit, and serum creatinine higher than 1.4 mg/dL (124  $\mu$ mol/L).

Also excluded were potential participants using systemic or inhaled corticosteroids or lithium, having a history of alcohol abuse within the past year, or having a history or presence of a sig-

nificant psychiatric disorder or any condition that, in the investigator's judgment, would interfere with participation in the trial. Potential participants who initiated a new drug therapy within 30 days of randomization, who were already participating in a weight loss program, or who took prescription weight loss or investigational medications within 90 days of randomization were also excluded. If clinically significant abnormalities were found at screening, individuals were referred to their personal physician. The protocol was reviewed and approved annually by the institutional review board at each site.

### Randomization


After all screening test results were reviewed and the candidate's eligibility confirmed, a randomization envelope prepared by the data coordinating center was opened and the participant was assigned to self-help or the commercial program (FIGURE 1). Participants were block randomized as determined by a random number table. A different block and randomization sequence was prepared for each site. Participants and investigators at each site were blind to the assignment condition until the envelope was opened.

### Treatments

Participants assigned to the self-help group received 20-minute consultations with a dietitian at the week 0 (baseline) and week 12 visits and were given publicly available printed material orienting them to dietary principles and exercise guidelines for safe weight loss.<sup>16,17</sup> Other resources such as public library materials, Web sites, and telephone numbers of health promotion organizations offering free weight control information were drawn to their attention.

Participants assigned to the commercial program were given vouchers entitling them to attendance at sessions of Weight Watchers, and the locations of available sites of this commercial program were reviewed with them. The vouchers enabled participants to attend sessions at no cost and, during the study

**Figure 1.** Flow of Participants in the Trial



BMI indicates body mass index.

period, had a retail value of approximately \$9 per voucher. The commercial program consists of a food plan, an activity plan, and a behavior modification plan focused primarily on cognitive restructuring. The food plan is a nutritionally balanced, moderate-deficit diet designed to produce a weight loss of up to 0.9 kg/wk. The activity plan follows current National Institutes of Health guidelines.<sup>18</sup> Weekly group meetings of approximately an hour's duration are led by successful program graduates who act as role models and provide written educational materials, a weekly weigh-in, and social support.

Compliance with treatment assignment in the commercial group was assessed at each clinic visit by self-reported attendance at group meetings. The use of concomitant medications for weight loss, including herbal preparations, was recorded.

### Outcome Measures

The primary outcome measure was change in body weight measured at each visit. Secondary measures to assess change in adiposity were BMI, waist circumference (at weeks 26, 52, and 104), and body fat measured by bioimpedance analysis (Tanita Bodyfat Analyzer, model TBF 105 or 305, Tanita Corporation, Arlington Heights, Ill). Other secondary measures were blood pressure (measured at each visit); total cholesterol, high-density lipoprotein (HDL) cholesterol, triglycerides, insulin, and glucose (measured at 1-year intervals); and quality of life measured using the Medical Outcomes Study Short-Form 36 Health Survey (SF-36)<sup>19</sup> and Impact of Weight on Quality of Life Questionnaire (IWQOL-Lite)<sup>20</sup> scales.

Adverse events, vital signs, and a standard panel of 20 biochemical indicators were monitored for safety. Blood samples were sent to a commercial laboratory (Quest Diagnostics, San Juan Capistrano, Calif) except for insulin lev-

els, which were assayed at the New York Obesity Research Center, New York City.

### Statistical Analysis

Differences between the treatment groups at randomization and characteristics of completers vs dropouts were tested using independent *t* tests for continuous variables and the Fisher exact test for categorical variables. Values are presented as mean (SD) except where otherwise noted. Statistical significance was set at  $P < .05$ .

Three analyses were performed on weight change and related outcome variables: an intent-to-treat (ITT) analysis including all randomized participants (missing values were imputed by last-observation-carried-forward or linear interpolation and participants who made no follow-up visits were assumed to remain at baseline value); a modified ITT analysis including all participants who made at least 1 clinic visit after randomization; and a completers

analysis using only participants who completed the study.

A mixed-model repeated-measures analysis of covariance was used to assess the significance of changes in the outcome variable, with treatment group, clinic site, visit, and visit  $\times$  group interaction as fixed effects, and initial weight as a covariate (Proc Mixed, SAS version 8, SAS Institute Inc, Cary, NC). Similar mixed regression models were used for analysis of changes in secondary outcome measures. Initial values of variables such as age, sex, and BMI were included as covariates if they contributed significantly to the model.

In the completers analysis of biological indices participants taking blood pressure, lipid, or glucose lowering medication were excluded from analyses on those measures, respectively, but included in all others. Analyses on changes in biological indices were run once including all values and a second time excluding values more than 3 SDs from the mean. Results were similar in all substantive aspects and the latter are presented here as being more typical of the interventions. The relation between weight changes and biological indicators of health was examined by including weight change and interactions of weight change and treatment group in linear regression models.

For the tabulation of cases by percentage of weight lost, separate  $\chi^2$  tests were conducted for year 1 and year 2 to test the hypothesis that the distributions in the self-help group were not different from those in the commercial group. Following rejection of the null hypothesis, frequencies within each percentage of weight loss category were tested against the null hypothesis that successes were distributed in the same proportion as the number of participants in the 2 groups.

## RESULTS

### Participant Characteristics

Characteristics of participants at baseline are shown in TABLE 1. Two participants developed lymphoma during the trial and were excluded from the completers analysis. Two participants assigned to the commercial group re-

**Table 1.** Baseline Characteristics of Participants by Randomization Group\*

| Variables                   | Mean (SD)                      |                               |
|-----------------------------|--------------------------------|-------------------------------|
|                             | Commercial Group<br>(n = 211)‡ | Self-help Group<br>(n = 212)§ |
| Age, y                      | 45 (10)                        | 44 (10)                       |
| Women, No. (%)              | 173 (82)                       | 185 (87)                      |
| Smokers, No. (%)            | 21 (10)                        | 19 (9)                        |
| Height, cm                  | 167 (8)                        | 166 (8)                       |
| Weight, kg                  | 94.2 (13.1)                    | 93.1 (14.4)                   |
| Body mass index†            | 33.8 (3.4)                     | 33.6 (3.7)                    |
| Fat, %‡                     | 44 (8)                         | 45 (9)                        |
| Waist, cm                   | 101 (12)                       | 99 (12)                       |
| Blood pressure, mm Hg       |                                |                               |
| Systolic                    | 123 (14)                       | 121 (12)                      |
| Diastolic                   | 79 (8)                         | 79 (9)                        |
| Glucose, mg/dL              | 92 (14)                        | 89 (12)                       |
| Insulin, IU/L               | 18 (9)                         | 18 (10)                       |
| Cholesterol, mg/dL          |                                |                               |
| Total                       | 214 (38)                       | 209 (36)                      |
| HDL                         | 50 (12)                        | 49 (13)                       |
| HDL/total cholesterol ratio | 0.24 (0.06)                    | 0.24 (0.06)                   |
| Triglycerides, mg/dL        | 155 (99)                       | 141 (82)                      |

Abbreviation: HDL, high-density lipoprotein.

SI conversion factors: To convert cholesterol to mmol/L, multiply values by 0.0259; to convert glucose to mmol/L, multiply values by 0.0555; to convert insulin to pmol/L, multiply values by 6.945; to convert triglycerides to mmol/L, multiply values by 0.0113.

\*There were no statistically significant differences between the commercial and self-help groups.

†Body mass index is measured as weight in kilograms divided by height in meters squared.

‡For fat %, n = 197, waist, n = 209, glucose and lipids, n = 210.

§For fat %, n = 194, waist, n = 209, triglycerides and insulin, n = 211.

**Table 2.** Change in Weight, Body Mass Index, and Waist Circumference by Analysis Method and Treatment Group

|                  | Change, Mean (SE)           |                            |                       |                                   |                            |                       |                             |                            |                       |
|------------------|-----------------------------|----------------------------|-----------------------|-----------------------------------|----------------------------|-----------------------|-----------------------------|----------------------------|-----------------------|
|                  | Intent-to-Treat Analysis    |                            |                       | Modified Intent-to-Treat Analysis |                            |                       | Completers Analysis         |                            |                       |
|                  | Commercial Group (n = 211)* | Self-help Group (n = 212)† | Between-Group P Value | Commercial Group (n = 198)*       | Self-help Group (n = 188)† | Between-Group P Value | Commercial Group (n = 148)* | Self-help Group (n = 159)† | Between-Group P Value |
| Weight, kg       |                             |                            |                       |                                   |                            |                       |                             |                            |                       |
| Year 1           | -4.3 (0.4)                  | -1.3 (0.4)                 | <.001                 | -4.5 (0.5)                        | -1.4 (0.5)                 | <.001                 | -5.0 (0.5)                  | -1.4 (0.5)                 | <.001                 |
| Year 2           | -2.9 (0.5)                  | -0.2 (0.4)                 | <.001                 | -2.7 (0.5)                        | -0.1 (0.5)                 | <.001                 | -3.0 (0.6)                  | -0.1 (0.6)                 | <.001                 |
| Body mass index‡ |                             |                            |                       |                                   |                            |                       |                             |                            |                       |
| Year 1           | -1.6 (0.2)                  | -0.5 (0.2)                 | <.001                 | -1.7 (0.2)                        | -0.6 (0.2)                 | <.001                 | -1.9 (0.2)                  | -0.6 (0.2)                 | <.001                 |
| Year 2           | -1.1 (0.2)                  | -0.2 (0.2)                 | <.001                 | -1.1 (0.2)                        | -0.1 (0.2)                 | <.001                 | -1.2 (0.2)                  | -0.1 (0.2)                 | <.001                 |
| Waist, cm§       |                             |                            |                       |                                   |                            |                       |                             |                            |                       |
| Year 1           | -4.1 (0.6)                  | -1.6 (0.6)                 | .003                  | -4.4 (0.7)                        | -1.8 (0.7)                 | .007                  | -4.9 (0.8)                  | -1.9 (0.8)                 | .005                  |
| Year 2           | -2.4 (0.6)                  | -0.6 (0.6)                 | .02                   | -2.3 (0.7)                        | -0.2 (0.7)                 | .03                   | -2.6 (0.7)                  | -0.2 (0.7)                 | .02                   |

\*For the commercial group, all values are different from baseline at  $P < .001$  for weight and body mass index. For change in waist circumference,  $P < .01$  vs baseline for year 1 and  $P < .04$  vs baseline for year 2.

†For the self-help group, all year 1 values are different from baseline at  $P < .01$  except for waist change under completers analysis ( $P < .02$ ); all year 2 values are not significantly different from baseline ( $P > .05$ ).

‡Body mass index is measured as weight in kilograms divided by height in meters squared.

§Because some waist measurements were not taken at baseline,  $n = 209$  in both groups for the intent-to-treat analysis and  $n = 146$  for commercial and  $n = 157$  for self-help for the completers analysis.


ported using a weight loss medication and were excluded from completers efficacy analyses. In the self-help group almost all participants reported attempting to change diet and increase physical activity, 14 reported using weight loss medications, another 6 tried herbal products, 10 enrolled in some form of structured commercial program (TOPS [Take Off Pounds Sensibly], Jenny Craig, 5 in Weight Watchers), and 9 mentioned following an alternative diet plan (protein, Atkins, the Zone) at some point during the 2-year study. All were retained in the analyses.

Participants who dropped out of the study were younger (age, 42.0 [10.6] vs 45.8 [9.9] years), had a higher BMI (34.3 [3.5] vs 33.4 [3.5]), greater percentage of body fat (46.6% [8.8%] vs 44.1% [8.3%]), were more likely to be smokers, and reported slightly lower income than participants who completed the study (for all comparisons  $P < .05$ ). The characteristics of dropouts did not differ by treatment group.

**Safety Parameters**

Bonferroni-adjusted statistical tests on the standard safety panel of 20 biochemistry analytes at randomization found only 1 difference: mean alanine aminotransferase values in the commercial group were statistically higher than in the self-help group, although

**Figure 2.** Mean Weight Changes by Group



Error bars indicate SEM.

both were within the normal range and the difference is not clinically significant (25.7 [13.8] vs 21.7 [11.2] U/L,  $P < .001$ ). There were no clinically significant changes in safety parameters over the course of the study in either group and no participants were removed from the study for adverse changes in biochemistry parameters.

**Weight, BMI, Body Fat, and Waist Circumference Outcomes**

Participants' change in weight ranged from -28 to +12 kg in the commercial group and -26 to +15 kg in the self-help group at 1 year, and -23 to +21

kg in the commercial group and -26 to +30 kg in the self-help group at the end of year 2.

Results of ITT, modified ITT, and completers analyses of weight changes and related body composition variables are shown in TABLE 2. Because of the low attrition rate, the results are not greatly different among these 3 analyses.

Mean weight changes from baseline for each of the clinic visits for all participants who made the visit are shown in FIGURE 2. At each visit, weight in the commercial group was significantly lower than at baseline and the amount of loss was greater than in the self-

help group. Weight in the self-help group was significantly less than baseline weight until week 52 but not thereafter. There were differences among sites in the amount of weight loss ( $P < .01$ ); however, the magnitude of the difference in weight loss between commercial and self-help groups at the 6 sites was similar (the treatment  $\times$  site interaction was not significant). The difference from baseline decreased over time in both groups after week 26 ( $P < .001$ ) and the rates of regain were not different (the treatment  $\times$  time interaction was not significant). Amount of weight lost by men compared with women was not significantly different, although the average weight of men

was greater at baseline (106.7 [11.9] vs 91.2 [12.7] kg;  $P < .001$ ). Weight loss and gain as a percentage of initial weight at 1 and 2 years is shown in TABLE 3.

Compliance with treatment assignment in the commercial group was assessed by self-reported attendance at meetings. The median number of sessions attended for 6-month periods ending at weeks 26, 52, 78, and 104 were 21, 19, 17, and 13, respectively. Weight loss differed as a function of self-reported attendance level ( $P < .05$ ) (FIGURE 3). Some weight regain was seen between year 1 and 2 at all attendance levels ( $P < .001$ ).

Analysis of changes in BMI yielded results similar to those for weight loss. Af-

ter 1 year 56% and after 2 years 52% of the commercial group participants were more than 1 BMI unit below starting BMI, compared with 31% after 1 year and 29% after 2 years in the self-help group. The differences between commercial and self-help groups in waist circumference reduction at 1 and 2 years were statistically significant (Table 2). The amount of reduction declined over time in both groups but a difference of about 2 cm remained at the end of 2 years. There were also significant differences between commercial and self-help groups in average amount of fat loss maintained over the last 18 months (3.2 [6.3] kg vs 1.6 [6.4] kg, respectively;  $P = .04$ ). Mean fat loss by men was not significantly different from that of women.

**Table 3.** Weight Change as a Percentage of Initial Weight, by Treatment Group\*

|                                 | Year 1†                    |                           |         | Year 2‡                    |                           |         |
|---------------------------------|----------------------------|---------------------------|---------|----------------------------|---------------------------|---------|
|                                 | No. (%)                    |                           | P Value | No. (%)                    |                           | P Value |
|                                 | Commercial Group (n = 172) | Self-help Group (n = 170) |         | Commercial Group (n = 148) | Self-help Group (n = 159) |         |
| Decrease from initial weight, % |                            |                           |         |                            |                           |         |
| ≥10                             | 37 (21)                    | 15 (9)                    | .002    | 24 (16)                    | 9 (6)                     | .005    |
| >5-<10                          | 29 (17)                    | 19 (11)                   | .15     | 27 (18)                    | 24 (15)                   | .57     |
| 0-5                             | 68 (39)                    | 57 (33)                   | .37     | 42 (28)                    | 46 (29)                   | >.99    |
| Increase from initial weight, % |                            |                           |         |                            |                           |         |
| 0-5                             | 32 (19)                    | 60 (35)                   | .003    | 38 (26)                    | 50 (31)                   | .41     |
| >5-<10                          | 5 (3)                      | 17 (10)                   | .01     | 15 (10)                    | 15 (9)                    | .71     |
| ≥10                             | 1 (1)                      | 2 (1)                     | ...     | 2 (1)                      | 15 (9)                    | .003    |

\*Percentages may not sum to 100% due to rounding. Ellipsis indicates numbers too small to perform statistical analysis.

† For year 1,  $\chi^2_{(df=2)} = 27.6$ ;  $P < .001$  for hypothesis of similar frequency distribution in the 2 groups.

‡ For year 2,  $\chi^2_{(df=2)} = 18.4$ ;  $P = .002$  for hypothesis of similar frequency distribution in the 2 groups.


**Biological Indices and Quality of Life Scales**

Although there were sustained differences in mean weight loss between the 2 groups over the duration of the study, the wide range in degree of individual success in each group and the low correlation between amount of weight loss and improvement in biological parameters resulted in mainly nonsignificant differences between treatment groups (TABLE 4). Two indices, diastolic blood pressure and serum insulin, were statistically improved in the commercial group compared with self-help at year 1 but only insulin was significantly different at year 2. Total cholesterol and the HDL/total cholesterol ratio improved in both groups, whereas fasting glucose levels increased in both groups.

Weight changes were reliably correlated with changes in all 8 biological indices in the commercial group and in 5 of the 8 indices in the self-help group (all  $P < .05$ ). Changes in waist circumference were also significantly related to changes in all indices (all  $P < .02$ ). Change in total cholesterol and triglycerides was 3 or more times as large per unit change in waist circumference for men as for women.

There were no significant differences between groups in amount of improvement on quality of life scales. Weight loss and percentage of weight

**Figure 3.** Median Weight Loss in the Commercial Group by Self-reported Attendance



loss were positively correlated with improvements in all scales of the SF-36 at both year 1 and year 2 in both groups (data not shown). Using a Bonferroni-adjusted significance level of  $P=.006$ , at 1 year weight loss was significantly correlated with improvements in scales for general physical health ( $r=0.25$ ) and vitality ( $r=0.26$ ). At year 2, scales for physical function ( $r=0.22$ ) and vitality ( $r=0.21$ ) were significantly correlated with weight loss. A similar pattern was seen for scales of the IWQOL-Lite (data not shown). Weight strongly predicted total score and all subscale scores, with the strongest relationships for public distress, physical function, and total score. The relationship was in the expected direction, with lower weight predicting better quality of life, and this did not differ by treatment.

#### COMMENT

The self-help group was able to lose and maintain approximately 1.3 to 1.4 kg for the first year, after which weight tended to increase until it returned to baseline at 2 years. The commercial group maintained a weight loss of 4.3 to 5.0 kg at the end of the first year and was 2.7 to 3.0 kg lower than baseline weight at the end of the second year. Participants who attended 78% or more of the commercial group sessions main-

tained a mean weight loss of almost 5 kg at the end of the 2-year study. These weight losses were achieved despite a general tendency to gain weight during most of the adult lifespan.<sup>21</sup> After 2 years, 52% of the commercial group and 29% of the self-help group had a BMI 1 unit or more below baseline BMI, an amount that the Institute of Medicine report on obesity treatment described as significant long-term weight loss.<sup>22</sup> Waist circumference, an independent cardiovascular risk factor,<sup>23,24</sup> was also reduced in the commercial group by about 4.5 cm at 1 year and 2.5 cm at 2 years. It is noteworthy that these results were obtained not in academic or research settings but in the context of a regular ongoing commercial program.

The correlations of changes in biological parameters with weight changes were generally similar to those seen in other studies with modest amounts of weight loss.<sup>4,25-29</sup> The magnitude of the group change may depend on characteristics of the study sample at randomization and the precise nature of the intervention (eg, diet, exercise, sodium content of diet, amount of weight lost). The reasons for the low correlations of individual improvement in biological indices with amount of weight lost are not known. Additional analyses are being conducted to establish whether

individual participant factors, such as the nature of the dietary and lifestyle changes, might account in part for the degree of improvement.

Despite sustained differences in mean weight loss between the 2 groups, the mean improvement in most biological parameters was not statistically different. In part, this is a statistical issue because the study was powered to detect differences in weight loss, not biological indices. However, the wide variability in attendance at the commercial sessions and success at weight loss highlights the importance of compliance with the assigned treatment. In this case, attendance at group meetings may provide the conditions necessary to produce and sustain weight loss and may increase the likelihood of improvements in biological indicators. However, even among those with the highest degree of adherence, there was weight regain during the second year of the study.

Self-selection bias—the tendency for successful participants to remain in the trial—may be a factor influencing these weight loss results, although that influence is likely to be relatively small because the high retention rate (73% after 2 years) limited the opportunity for self-selection to occur. Also, the correlation between attendance and success at weight loss is supported by other

**Table 4.** Changes in Biological Indices by Year and Treatment Group

|                             | Change, Mean (SE) |                 |                       |                  |                 |                       |
|-----------------------------|-------------------|-----------------|-----------------------|------------------|-----------------|-----------------------|
|                             | Year 1            |                 |                       | Year 2           |                 |                       |
|                             | Commercial Group  | Self-help Group | Between-Group P Value | Commercial Group | Self-help Group | Between-Group P Value |
| Blood pressure, mm Hg       |                   |                 |                       |                  |                 |                       |
| Systolic                    | -0.6 (0.9)        | 0.2 (0.8)       | .53                   | -2.2 (1.1)*      | -2.4 (1.0)*     | .93                   |
| Diastolic                   | -0.4 (0.6)        | 1.4 (0.6)*      | .02                   | -0.6 (0.7)       | 0 (0.6)         | .51                   |
| Glucose, mg/dL              | 3.5 (0.6)*        | 3.6 (0.6)*      | .90                   | 5.2 (0.7)*       | 4.6 (0.7)*      | .47                   |
| Insulin, IU/L               | -2.0 (0.5)*       | -0.3 (0.5)      | .01                   | 0.6 (0.6)        | 2.3 (0.6)*      | .04                   |
| Cholesterol, mg/dL          |                   |                 |                       |                  |                 |                       |
| Total                       | -8.7 (1.7)*       | -9.5 (1.7)*     | .73                   | -10.4 (2.0)*     | -11.3 (1.9)*    | .74                   |
| HDL                         | 2.0 (0.7)*        | 0.8 (0.7)       | .17                   | 0.5 (0.8)        | 0.6 (0.8)       | .92                   |
| HDL/total cholesterol ratio | 0.02 (0.003)*     | 0.020 (0.003)*  | .43                   | 0.02 (0.003)*    | 0.02 (0.003)*   | .99                   |
| Triglycerides, mg/dL        | -7.8 (3.7)*       | 1.5 (3.8)       | .08                   | -0.3 (4.0)       | -0.1 (3.9)      | .97                   |

Abbreviation: HDL, high-density lipoprotein.

SI conversion factors: To convert cholesterol to mmol/L, multiply values by 0.0259; to convert glucose to mmol/L, multiply values by 0.0555; to convert insulin to pmol/L, multiply values by 6.945; to convert triglycerides to mmol/L, multiply values by 0.0113.

\*Value is significantly different from baseline at  $P<.05$ .

research showing that length of the treatment period, and consequently the number of intervention meetings, has an independent effect on success.<sup>30</sup> It is also possible that characteristics of individuals willing and motivated to participate in long-term randomized clinical trials may have influenced the trial results. Finally, it should be emphasized that this study reports on a particular commercial program with many unique aspects. Our results should not be taken as representative of all commercial programs, many of which use other interventions, such as proprietary liquid formulas or diets that are not balanced.

In summary, this 2-year trial provides information on weight loss in an ongoing structured commercial weight loss program in comparison with self-help attempts to lose weight. The results show that this program

provides modest weight loss but is more effective than brief counseling and self-help for overweight and obese adults.

**Author Contributions:** As principal investigator of the coordinating center, Dr Heshka had full access to all the data in this study and takes responsibility for the integrity of the data and the accuracy of the data analysis. **Study concept and design:** Heshka, Anderson, Atkinson, Greenway, Hill, Phinney, Miller-Kovach, Pi-Sunyer. **Acquisition of data:** Anderson, Atkinson, Greenway, Hill, Phinney, Kolotkin, Pi-Sunyer. **Analysis and interpretation of data:** Heshka, Anderson, Atkinson, Hill, Kolotkin, Miller-Kovach, Pi-Sunyer. **Drafting of the manuscript:** Heshka, Anderson, Hill, Kolotkin. **Critical revision of the manuscript for important intellectual content:** Heshka, Anderson, Atkinson, Greenway, Hill, Phinney, Miller-Kovach, Pi-Sunyer. **Statistical expertise:** Heshka. **Obtained funding:** Anderson, Miller-Kovach, Pi-Sunyer. **Administrative, technical, or material support:** Anderson, Atkinson, Greenway, Hill, Phinney, Kolotkin, Miller-Kovach, Pi-Sunyer. **Study supervision:** Heshka, Anderson, Greenway, Hill, Phinney, Pi-Sunyer. **Funding/Support:** This study was supported by a grant from Weight Watchers International (Woodbury, NY) to the New York Obesity Research Center at St Luke's/Roosevelt Hospital.

**Role of the Sponsor:** The sponsor, Weight Watchers International, initiated the project by contacting the New York Obesity Research Center and offering to fund a rigorous long-term study to evaluate the success of its program. The study design evolved during a series of discussions with the sponsor and coinvestigators and included a 1-day meeting of all the principal investigators, the Weight Watchers vice president for research and development (Ms Miller-Kovach), and the medical director of Weight Watchers to review and finalize the protocol. Funding was provided through a research grant to the coordinating center (New York Obesity Research Center at St Luke's/Roosevelt Hospital), which subsequently disbursed funds to the collaborating clinical sites and central laboratory. Data collected at the clinical sites and from the central laboratory were forwarded electronically to the coordinating center where they were pooled for analysis. The sponsor had no role in data managements or analysis and did not write any part of the manuscript. Questions that arose during the preparation of manuscripts were discussed among the appropriate coinvestigators, including the coinvestigator who is an employee of the sponsor. The research contract stipulates that the results of the trial may be published as a scientific paper and that the sponsor would be given opportunity to comment on the manuscript prior to submission for publication.

**Acknowledgment:** We gratefully acknowledge the contribution and assistance of the study coordinators and medical staff at each of the clinical study sites and of Myron Winick, MD, Weight Watchers Medical Director.

## REFERENCES

- Serdula MK, Mokdad AH, Williamson DF, Galuska DA, Mendlein JM, Heath GW. Prevalence of attempting weight loss and strategies for controlling weight. *JAMA*. 1999;282:1353-1358.
- Serdula MK, Collins ME, Williamson DF, Anda RF, Pamuk E, Byers TE. Weight control practices of US adolescents and adults. *Ann Intern Med*. 1993;119:667-671.
- Levy AS, Heaton AW. Weight control practices of US adults trying to lose weight. *Ann Intern Med*. 1993;119:661-666.
- National Heart, Lung, and Blood Institute. *Clinical Guidelines on the Identification, Evaluation, and Treatment of Overweight and Obesity in Adults: The Evidence Report*. Bethesda, Md: National Institutes of Health; June 1998.
- Khaodhlar L, Blackburn GL. Health benefits and risks of weight loss. In: Bjorntorp P, ed. *International Textbook of Obesity*. Chichester, England: John Wiley & Sons; 2001:413-439.
- Klein S. Outcome success in obesity. *Obes Res*. 2001;9:3545-3585.
- Goldstein DJ. Beneficial health effects of modest weight loss. *Int J Obes Relat Metab Disord*. 1992;16:397-415.
- Academy Advisory Group, National Academy of Sciences. *Preventive Services for a Well Population*. Washington, DC: National Academy of Sciences; 1978.
- Rippe JM, Price JM, Hess SA, et al. Improved psychological well-being, quality of life and health practices in moderately overweight women participating in a 12-week structured weight loss program. *Obes Res*. 1998;6:208-218.
- Lowe MR, Miller-Kovach K, Frye N, Phelan S. An initial evaluation of a commercial weight loss program: short-term effects on weight, eating behavior, and mood. *Obes Res*. 1999;7:51-59.
- Taylor CB, Stunkard AJ. Public health approaches to weight control. In: Stunkard AJ, Wadden TA, eds. *Obesity: Theory and Therapy*. 2nd ed. New York, NY: Raven Press; 1993:335-353.
- Wadden TA, Foster GD, Letizia KA, Stunkard AJ. A multicenter evaluation of a proprietary weight reduction program for the treatment of marked obesity. *Arch Intern Med*. 1992;152:961-966.
- Lowe MR, Miller-Kovach K, Phelan S. Weight-loss maintenance in overweight individuals one to five years following successful completion of a commercial weight loss program. *Int J Obes Relat Metab Disord*. 2001;25:325-331.
- The US Weight Loss and Diet Control Market*. Tampa, Fla: Marketdata Enterprises Inc; 2001.
- Heshka S, Greenway F, Anderson JW, et al. Self-help weight loss versus a structured commercial program after 26 weeks: a randomized controlled study. *Am J Med*. 2000;109:282-287.
- Eating Smart*. Chicago, Ill: National Cattlemen's Beef Association; 1996. Distributed by Shape Up America.
- Fitting Fitness In*. Chicago, Ill: National Cattlemen's Beef Association; 1996. Distributed by Shape Up America.
- Physical Activity and Cardiovascular Health*. Bethesda, Md: National Institutes of Health; 1995. NIH Consensus Statement.
- Ware JE Jr. *SF-36 Health Survey: Manual & Interpretation Guide*. Boston, Mass: The Health Institute; 1997.
- Kolotkin RL, Crosby RD, Kosloski KD, Williams GR. Development of a brief measure to assess quality of life in obesity. *Obes Res*. 2001;9:102-111.
- Heo M, Faith MS, Mott JW, Gorman BS, Redden DT, Allison DB. Tutorial on hierarchical linear models for the development of growth curves: an example with body mass index in overweight/obese adults. *Stat Med*. In press.
- Thomas PR, ed, and the Food and Nutrition Board, Institute of Medicine, Committee to Develop Criteria for Evaluating the Outcomes of Approaches to Prevent and Treat Obesity. *Weighing the Options: Criteria for Evaluating Weight Management Programs*. Washington, DC: National Academy Press; 1995.
- Kannel WB, Cupples LA, Ramaswami R, et al. Regional obesity and risk of cardiovascular disease: the Framingham Study. *J Clin Epidemiol*. 1991;44:183-190.
- Janssen I, Katzmarzyk PT, Ross R. Body mass index, waist circumference, and health risk: evidence in support of current National Institutes of Health guidelines. *Arch Intern Med*. 2002;162:2074-2079.
- Stevens VJ, Obarzanek E, Cook NR, et al. Long-term weight loss and changes in blood pressure: results of the trials of hypertension prevention, phase II. *Ann Intern Med*. 2001;134:1-11.
- Ashley JM, Tanaka-St Jeor S, Perumean-Chaney S, Schrage J, Bovee V. Meal replacements in weight intervention. *Obes Res*. 2001;9(suppl 4):3125-3205.
- Dattilo AM, Kris-Etherton PM. Effects of weight reduction on blood lipids and lipoproteins: a meta-analysis. *Am J Clin Nutr*. 1992;56:320-328.
- Anderson JW, Konz EC. Obesity and disease: effects of weight loss on comorbid conditions. *Obes Res*. 2001;9(suppl 4):3265-3345.
- Mertens IL, Van Gaal LF. Overweight, obesity and blood pressure: the effects of modest weight reduction—a review. *Obes Res*. 2000;8:270-278.
- Perri MG, Nezu AM, Patti ET, McCann KL. Effect of length of treatment on weight loss. *J Consult Clin Psychol*. 1989;57:450-452.