



# Outbreaks of Avian Influenza A (H5N1) in Asia and Interim Recommendations for Evaluation and Reporting of Suspected Cases—United States, 2004

Online article and related content current as of November 26, 2009.

JAMA. 2004;291(10):1191-1193 (doi:10.1001/jama.291.10.1191)

<http://jama.ama-assn.org/cgi/content/full/291/10/1191>

Correction

[Contact me if this article is corrected.](#)

Citations

[Contact me when this article is cited.](#)

Topic collections

Viral Infections; Infectious Diseases  
[Contact me when new articles are published in these topic areas.](#)

Subscribe

<http://jama.com/subscribe>

Permissions

[permissions@ama-assn.org](mailto:permissions@ama-assn.org)  
<http://pubs.ama-assn.org/misc/permissions.dtl>

Email Alerts

<http://jamaarchives.com/alerts>

Reprints/E-prints

[reprints@ama-assn.org](mailto:reprints@ama-assn.org)



## Outbreaks of Avian Influenza A (H5N1) in Asia and Interim Recommendations for Evaluation and Reporting of Suspected Cases—United States, 2004

MMWR. 2004;53:97-100

DURING DECEMBER 2003–FEBRUARY 2004, outbreaks of highly pathogenic avian influenza A (H5N1) among poultry were reported in Cambodia, China, Indonesia, Japan, Laos, South Korea, Thailand, and Vietnam. As of February 9, 2004, a total of 23 cases of laboratory-confirmed influenza A (H5N1) virus infections in humans, resulting in 18 deaths, had been reported in Thailand and Vietnam. In addition, approximately 100 suspected cases in humans are under investigation by national health authorities in Thailand and Vietnam. CDC, the World Health Organization (WHO), and national health authorities in Asian countries are working to assess and monitor the situation, provide epidemiologic and laboratory support, and assist with control efforts. This report summarizes information about the human infections and avian outbreaks in Asia and provides recommendations to guide influenza A (H5N1) surveillance, diagnosis, and testing in the United States.

### Poultry Outbreaks

On December 12, 2003, an outbreak of avian influenza A (H5N1) among poultry in South Korea was reported. Subsequent influenza A (H5N1) outbreaks among poultry were confirmed in Vietnam (January 8, 2004), on a single farm in Japan (January 12), in

Thailand (January 23), in Cambodia (January 24), in China (January 27), in Laos (January 27), and in Indonesia (February 2). On January 19, a single peregrine falcon found dead in Hong Kong also tested positive for influenza A (H5N1) virus, but no poultry outbreak has been identified.

In Vietnam, as of February 9, a total of 18 human influenza A (H5N1) infections had been reported, resulting in 13 deaths. Patients ranged in age from 4 to 30 years; 10 patients were aged <18 years. The cases included fatal infections in two sisters who were part of a cluster of four cases of severe respiratory illness in a single family.

In Thailand, influenza A (H5N1) infection was confirmed in four males, aged 6-7 years, and one female, aged 58 years. All five patients died.<sup>1</sup> Other cases are under investigation.

### Analysis of Viruses

Antigenic analysis and genetic sequencing distinguish between influenza viruses that usually circulate among birds and those that usually circulate among humans. Sequencing of the H5N1 viruses obtained from five persons in Vietnam and Thailand, including one sister from the cluster in Vietnam, has indicated that all of the genes of these viruses are of avian origin. No evidence of genetic reassortment between avian and human influenza viruses has been identified. If reassortment occurs, the likelihood that the H5N1 virus can be transmitted more readily from person to person will increase. Although all the genes are of avian origin, the current H5N1 viruses are antigenically distinguishable from those isolated from humans in Hong Kong in 1997 and 2003.

Genetic sequencing of the five human H5N1 isolates from Thailand and Vietnam also indicates that the viruses have genetic characteristics associated with resistance to the influenza antiviral drugs amantadine and

rimantadine. Antiviral susceptibility testing confirms this finding. Testing for susceptibility of the H5N1 isolates to the neuraminidase inhibitor oseltamivir has demonstrated the sensitivity of these viruses to the drug; testing to determine susceptibility to the neuraminidase inhibitor zanamavir is under way.

### Interim Recommendations for U.S. Surveillance and Diagnostic Evaluation

CDC recommends that state and local health departments, hospitals, and clinicians enhance their efforts to identify patients who could be infected by influenza A (H5N1) virus and take infection-control precautions when influenza A (H5N1) is suspected (see sidebar). Testing of hospitalized patients for influenza A (H5N1) infection is indicated when both of the following exist: (1) radiographically confirmed pneumonia, acute respiratory distress syndrome (ARDS), or other severe respiratory illness for which an alternative diagnosis has not been established and (2) a history of travel within 10 days of symptom onset to a country with documented H5N1 avian influenza infections in poultry or humans. Ongoing listings of countries affected by avian influenza are available from the World Organization for Animal Health.\*

Testing for influenza A (H5N1) also should be considered on a case-by-case basis in consultation with state and local health departments for hospitalized or ambulatory patients with all of the following: (1) documented temperature of >100.4°F (>38°C); (2) cough, sore throat, or shortness of breath; and (3) history of contact with poultry or domestic birds (e.g., visited a poultry farm, a household raising poultry, or a bird market) or a known or suspected patient with influenza A (H5N1) in an H5N1-affected country within 10 days of symptom onset.

**BOX. Interim recommended infection-control precautions\* for influenza A (H5N1)**

- All patients with a febrile respiratory illness should be asked about their recent travel history and managed using *Respiratory Hygiene/Cough Etiquette in Health Care Settings* guidelines<sup>†</sup>.
- Isolation precautions for all hospitalized patients who have or are under evaluation for influenza A (H5N1) are the same as those that should be used for severe acute respiratory syndrome (SARS), as follows:
  - Pay careful attention to hand hygiene before and after all patient contact.
  - Use gloves and gown for all patient contact.
  - Wear eye protection when within 3 feet of the patient.
  - Place the patient in an airborne isolation room (i.e., monitored negative air pressure in relation to surrounding areas with six to 12 air changes per hour).
  - When entering the patient's room, use a fit-tested respirator at least as protective as an N95 filtering-facepiece respirator approved by the National Institute for Occupational Safety and Health.
- Outpatients or hospitalized patients discharged in <14 days should be isolated in the home setting on the basis of principles for home isolation of SARS patients<sup>§</sup>.
- These precautions should be continued for 14 days after onset of symptoms until an alternative diagnosis is established or diagnostic test results indicate that the patient is not infected with influenza A virus.

\* Additional information about health-care isolation precautions is available at <http://www.cdc.gov/ncidod/hip/isolat/isolat.htm>.

† Available at [http://www.cdc.gov/flu/professionals/infectioncontrol/resp\\_hygiene.htm](http://www.cdc.gov/flu/professionals/infectioncontrol/resp_hygiene.htm).

§ Available at <http://www.cdc.gov/ncidod/sars/guidance>.

**Recommended Laboratory Testing Procedures**

The highly pathogenic avian influenza A (H5N1) virus requires Biosafety Level (BSL)-3+laboratory conditions for certain procedures. CDC recommends that virus isolation studies on respiratory specimens from patients who meet the testing criteria should not be performed unless all BSL-3+ conditions are met. However, clinical specimens can be tested by polymerase chain reaction (PCR) assays by using standard BSL-2 work practices in a Class II biological safety cabinet. CDC has developed real-time PCR protocols<sup>†</sup> for various respiratory pathogens, including SARS and influenza A and B viruses. In addition, commer-

cially available antigen-detection tests can be used under BSL-2 levels to test for influenza. Although these rapid tests for human influenza also can detect avian influenza A (H5N1) viruses, the sensitivity of these tests is substantially lower than that of virus culture or PCR.<sup>2</sup>

Specimens from persons meeting clinical and epidemiologic indications for testing should be sent to CDC if they test positive for influenza A either by PCR or antigen detection testing, or if PCR assays for influenza are not available locally. CDC also will accept, for follow-up testing, specimens from persons meeting the clinical and epidemiologic indications but testing negative on the rapid tests when PCR assay

was not available. Requests for testing by CDC should come through local and state health departments, which should contact CDC's Emergency Operations Center, telephone 770-488-7100.

Reported by: CDC/WHO Avian Influenza Response Team.

**CDC Editorial Note:** Since 1997, human infection with avian influenza viruses has been confirmed on five occasions.<sup>‡</sup> The ability of avian viruses to transmit from person to person appears limited. Rare person-to-person infection was noted in the A (H5N1) outbreak in Hong Kong in 1997<sup>3,4</sup> and in the A (H7N7) outbreak in the Netherlands in 2003,<sup>5</sup> but these secondary cases did not result in sustained chains of transmission or communitywide outbreaks. These previous experiences with avian influenza viruses suggest that limited person-to-person transmission of the current H5N1 viruses could occur.

The majority of the human H5N1 cases are apparently associated with direct exposure to infected birds or to surfaces contaminated with excretions from infected birds. The family respiratory illness cluster in Vietnam suggests the possibility of limited person-to-person transmission. However, other possibilities (e.g., transmission through exposure to surfaces contaminated by H5N1-infected poultry feces) cannot be ruled out. Although no evidence for sustained person-to-person transmission of influenza A (H5N1) has been identified, influenza viruses have the capacity to change quickly. Continued monitoring for new transmission patterns is an important aspect of the current investigation.

In 1997, the influenza A (H5N1) outbreak among persons in Hong Kong ended abruptly after the culling of poultry. However, the current outbreaks present challenges because of the large geographic areas and numbers of affected poultry. Asian poultry populations are maintained both on large commercial farms and in backyard flocks. In addition, infections among wild bird populations might be extensive, and the

resources to address this problem are limited in certain affected countries. Because of increasing evidence that avian influenza viruses infect humans, persons involved in the slaughter of poultry potentially infected with avian influenza viruses or their contaminated environments should follow WHO recommendations for worker protection.<sup>§</sup>

Because the influenza A (H5N1) virus could develop the ability to maintain sustained person-to-person transmission, WHO collaborating centers are working to coordinate vaccine development. Efforts are under way in the United Kingdom and the United States to develop influenza A (H5N1) reference viruses for use in vaccine preparation. The minimum estimated time necessary to complete reference virus development and safety testing is 3 months. Production by vaccine manufacturers of pilot lots of vaccine for clinical testing can begin only after reference virus development and safety testing have been completed. Decisions on whether to proceed with vaccine manufacture will depend, in part, on the evolution of the current outbreaks.

On February 4, CDC issued an order for an immediate ban<sup>||</sup> on the import of all birds from Cambodia, China (including Hong Kong), Indonesia, Japan, Laos, South Korea, Thailand, and Vietnam. Birds from these affected countries potentially can infect humans with influenza A (H5N1). This order complements a similar action taken by the U.S. Department of Agriculture (USDA).

CDC advises that travelers to countries in Asia with documented H5N1 outbreaks should avoid poultry farms, contact with animals in live food markets, and any surfaces that appear to be contaminated with feces from poultry or other animals. More information on travel is available from CDC at <http://www.cdc.gov/travel>. Additional information on influenza viruses and avian influenza is available from CDC at <http://www.cdc.gov/flu>. Updated information on hu-

man infections is available from WHO at <http://www.who.int/en>.

#### REFERENCES

1. CDC. Cases of influenza A (H5N1)—Thailand, 2004. *MMWR* 2004; 53:100-3.
2. CDC. Prevention and control of influenza: recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR* 2003; 52(No. RR-8).
3. Bridges CB, Lim W, Hu-Primmer J, et al. Risk of influenza A (H5N1) infection among poultry workers, Hong Kong, 1997-1998. *J Infect Dis* 2002; 185:1005-10.
4. Bridges CB, Katz JM, Seto WH, et al. Risk of influenza A (H5N1) infection among health care workers exposed to patients with influenza A (H5N1), Hong Kong. *J Infect Dis* 2000; 181:344-8.
5. de Jong JC, Rimmelzwaan GF, Bartelds AI, Wilbrink B, Fouchier RA, Osterhaus AD. The 2002/2003 influenza season in the Netherlands and the vaccine composition for the 2003/2004 season [Dutch]. *Ned Tijdschr Geneesk* 2003; 147:1971-5.

\*Available at [http://www.oie.int/eng/en\\_index.htm](http://www.oie.int/eng/en_index.htm).

†These protocols are available to public health laboratories and have been posted, under SARS (password required), by the Association of Public Health Laboratories at [http://www.aphl.org/members\\_only/index.cfm](http://www.aphl.org/members_only/index.cfm).

‡Influenza A (H5N1) in Hong Kong in 1997 and 2003, influenza A (H9N2) in Hong Kong in 1999 and 2003, and influenza A (H7N7) in the Netherlands in 2003.

§Available at <http://www.wpro.who.int/avian/docs/recommendations.asp>.

||Additional information on the embargo is available at <http://www.cdc.gov/flu/avian/embargo.htm>.

## Trends in Intake of Energy and Macronutrients—United States, 1971-2000

*MMWR*. 2004;53:80-82

1 table, 2 figures omitted

DURING 1971-2000, THE PREVALENCE OF obesity in the United States increased from 14.5% to 30.9%.<sup>1</sup> Unhealthy diets and sedentary behaviors have been identified as the primary causes of deaths attributable to obesity.<sup>2</sup> Evaluating trends in dietary intake is an important step in understanding the factors that contribute to the increase in obesity. To assess trends in intake of energy (i.e., kilocalories [kcal]), protein, carbohydrate, total fat, and saturated fat during 1971-2000, CDC analyzed data from four National Health and Nutri-

tion Examination Surveys (NHANES): NHANES I (conducted during 1971-1974), NHANES II (1976-1980), NHANES III (1988-1994), and NHANES 1999-2000. This report summarizes the results of that analysis, which indicate that, during 1971-2000, mean energy intake in kcals increased, mean percentage of kcals from carbohydrate increased, and mean percentage of kcals from total fat and saturated fat decreased (Figures 1 and 2). An expert advisory committee appointed by the U.S. Department of Health and Human Services and the U.S. Department of Agriculture (USDA) is conducting a review of the *Dietary Guidelines for Americans*.<sup>3</sup> Revised guidelines will be published in 2005.

NHANES provides information on the health and nutritional status of the U.S. civilian, noninstitutionalized population by using a complex, multi-stage probability sample design. NHANES I sampled persons residing in the contiguous 48 states; subsequent surveys sampled all 50 states. Surveys consisted of a household interview followed by an examination at a mobile examination center (MEC). All of the surveys included a dietary recall interview that was conducted at the MEC to obtain information on foods and beverages consumed during the preceding 24 hours. In this report, estimates of energy intake include kcals from alcoholic beverages; however, the percentage of kcals from alcohol is not presented separately. Age was recorded at the time of the household interview. The upper age limit was 74 years for NHANES I and NHANES II. No upper age limit was established for NHANES III and NHANES 1999-2000. To compare estimates across surveys, the analysis included only adults aged 20-74 years. Sample sizes ranged from 1,730 men and 2,003 women in NHANES 1999-2000 to 6,630 men and 7,537 women in NHANES III (Table).

Statistical analyses were conducted by using SAS version 8.2 and SUDAAN version 8.0.0, which used sample