



Opioid Abuse

Janet M. Torpy; Cassio Lynm; Richard M. Glass

JAMA. 2004;292(11):1394 (doi:10.1001/jama.292.11.1394)

Online article and related content
current as of November 8, 2009.

<http://jama.ama-assn.org/cgi/content/full/292/11/1394>

Supplementary material

Spanish PDF

<http://jama.ama-assn.org/cgi/content/full/292/11/1394/DC1>

Correction

[Contact me if this article is corrected.](#)

Citations

[Contact me when this article is cited.](#)

Topic collections

Public Health; Substance Abuse/ Alcoholism; JAMA Patient Page

[Contact me when new articles are published in these topic areas.](#)

Related Articles published in
the same issue

A 35-Year-Old Physician With Opioid Dependence

John R. Knight. *JAMA*. 2004;292(11):1351.

Subscribe

<http://jama.com/subscribe>

Email Alerts

<http://jamaarchives.com/alerts>

Permissions

permissions@ama-assn.org

<http://pubs.ama-assn.org/misc/permissions.dtl>

Reprints/E-prints

reprints@ama-assn.org

Opioid Abuse

Opioids are a family of related drugs that relieve pain. All of the opioids (sometimes called **narcotics**) are chemically related to opium, which is a substance collected from the poppy plant. Opioid drugs include opium, codeine, fentanyl, heroin, hydrocodone, methadone, morphine, oxycodone, paregoric, and sufentanil. When prescribed by a doctor, the pain-relieving properties of opioids are used during and after surgical procedures, for the pain of childbirth, for injury, and for other pain problems. Although opioid medications have helped millions of individuals with pain, these drugs can be used inappropriately. The September 15, 2004, issue of *JAMA* includes an article about abuse of prescription opioid medication.

ABUSE AND DEPENDENCE

Drug abuse is a pattern of inappropriate drug use that leads to recurrent problems in fulfilling obligations, impaired physical functioning, conflicts with family and friends, and legal problems. Drug abuse may progress to **dependence** (sometimes called **addiction**), manifested by a strong desire to continue the drug despite the increasingly severe problems it causes, **tolerance** (a need for larger amounts of the drug to get the same effects), and withdrawal symptoms if the drug is stopped.

SYMPTOMS AND SIGNS OF OPIOID ABUSE

- Sedation, sleepiness, or lethargy
- Avoidance and withdrawal from usual activities
- Multiple visits to multiple doctors to increase amounts of prescription drugs available for abuse
- Confusion
- Weight loss

COMPLICATIONS OF OPIOID ABUSE

- Blood-borne infections from unsterile injections, including human immunodeficiency virus (HIV), hepatitis viruses, and bacterial infections
- Job loss with possible financial devastation
- Loss of family, friends, and career-related relationships
- Increased chance of risky behavior, including driving under the influence
- Drug overdoses, which can lead to brain damage or death
- Miscarriage, stillbirth, or infants with low birth weight due to opioid abuse during pregnancy. Babies born to addicted mothers will be born addicted to the opioid and will have withdrawal symptoms after birth.

TREATMENT OF OPIOID ADDICTION

Recognition and admission that drug abuse exists is the first step in treatment. Drug addiction is a chronic medical problem. It is a treatable disease but relapse is a prominent feature. Relapse must be considered as part of a treatment plan. Drug abuse counseling is an important part of treatment. Participation in support groups, such as Narcotics Anonymous, may help individuals in treatment for opioid abuse. Certain medications may be used as a part of treatment. These include methadone, a long-acting opioid taken by mouth, which can substitute for the harmful injection of illegally obtained opioids.

FOR MORE INFORMATION

- National Institute on Drug Abuse
www.nida.nih.gov
- National Council on Alcoholism and Drug Dependence
800/622-2255
www.ncadd.org
- American Psychiatric Association
www.psych.org

INFORM YOURSELF

To find this and previous JAMA Patient Pages, go to the Patient Page link on *JAMA*'s Web site at www.jama.com. Many are available in English and Spanish. A Patient Page on cocaine addiction was published in the January 2, 2002, issue, and one on treating drug dependency was published in the March 8, 2000, issue.

Sources: National Institute on Drug Abuse, American Psychiatric Association, National Council on Alcoholism and Drug Dependence

Janet M. Torpy, MD, Writer

Cassio Lynn, MA, Illustrator

Richard M. Glass, MD, Editor

The JAMA Patient Page is a public service of *JAMA*. The information and recommendations appearing on this page are appropriate in most instances, but they are not a substitute for medical diagnosis. For specific information concerning your personal medical condition, *JAMA* suggests that you consult your physician. This page may be photocopied noncommercially by physicians and other health care professionals to share with patients. Any other print or online reproduction is subject to AMA approval. To purchase bulk reprints, call 718/946-7424.

