



Liver Transplantation

Janet M. Torpy; Cassio Lynm; Richard M. Glass

JAMA. 2006;295(18):2208 (doi:10.1001/jama.295.18.2208)

Online article and related content
current as of November 8, 2009.

<http://jama.ama-assn.org/cgi/content/full/295/18/2208>

Supplementary material

Spanish PDF

<http://jama.ama-assn.org/cgi/content/full/295/18/2208/DC1>

Correction

[Contact me if this article is corrected.](#)

Citations

[Contact me when this article is cited.](#)

Topic collections

Transplantation; Liver Transplantation; JAMA Patient Page;
Gastroenterology; Liver/ Biliary Tract/ Pancreatic Diseases
[Contact me when new articles are published in these topic areas.](#)

Related Articles published in
the same issue

Integrating Palliative Care for Liver Transplant Candidates: "Too Well for
Transplant, Too Sick for Life"
[Anne M. Larson et al. *JAMA*. 2006;295\(18\):2168.](#)

Subscribe

<http://jama.com/subscribe>

Email Alerts

<http://jamaarchives.com/alerts>

Permissions

permissions@ama-assn.org


<http://pubs.ama-assn.org/misc/permissions.dtl>

Reprints/E-prints

reprints@ama-assn.org

Liver Transplantation

The liver has many functions, including building proteins and other substances for the body to use, removal of waste products and toxins from the blood, drug metabolism, and energy storage. Liver disease causes these crucial functions to fail. When liver failure is too severe to respond to medications, liver transplantation may be an option. Because transplantation is a major surgical procedure and there is a shortage of available donor organs, not every person with liver failure can receive a transplant. At each transplant center, a multidisciplinary team reviews and selects patients who are then placed on a waiting list for a suitable donor organ. These donor organs are allocated by the organ donor coordinators based on need (most severely ill persons) and compatibility of donor and recipient. The May 10, 2006, issue of JAMA includes an article about serious liver disease and liver transplantation.



REASONS FOR LIVER TRANSPLANTATION

- Hepatitis C and hepatitis B—forms of hepatitis (liver infections) that can cause serious, permanent liver damage
- Cirrhosis (permanent liver damage from diseases such as primary biliary cirrhosis and inherited liver diseases or from alcohol abuse)
- In children, **biliary atresia** (failure of bile duct formation) is the most common reason for needing a liver transplant
- Toxic hepatic failure (can occur from eating poisonous mushrooms or from overdoses of drugs such as acetaminophen)
- **Hepatocellular carcinoma** (a primary cancer of the liver tissue), when found at an early, treatable stage

AFTER A LIVER TRANSPLANT

Liver transplantation is a major operation that takes place only in specialized transplant centers. The diseased liver is removed through an abdominal incision. The liver's blood supply (**portal vein**, **inferior vena cava**, and **hepatic artery**) and the **biliary system** are all connected to the new liver after it is placed into the abdominal cavity. Persons who have liver transplants require intensive care and close monitoring after their operation. In a few cases, only part of a healthy liver from a living (usually related) donor may be used. This requires the donor to have an operation to remove a part of their liver, which is then transplanted into the person with liver failure.

POSSIBLE TREATMENTS

- Immunosuppressive medications (drugs that prevent rejection of the transplanted organ) must be taken for the rest of the individual's life.
- The transplanted liver is monitored for function and for rejection, and the medication regimen is often adjusted.
- Persons with transplanted organs should avoid contact with ill individuals to prevent infection. Illnesses that healthy persons can tolerate have serious consequences for patients who are taking immunosuppressive medications.

Sources: National Institute of Diabetes and Digestive and Kidney Diseases; American Liver Foundation; United Network for Organ Sharing; American Gastroenterological Association

Janet M. Torpy, MD, Writer

Cassio Lynn, MA, Illustrator

Richard M. Glass, MD, Editor

The JAMA Patient Page is a public service of JAMA. The information and recommendations appearing on this page are appropriate in most instances, but they are not a substitute for medical diagnosis. For specific information concerning your personal medical condition, JAMA suggests that you consult your physician. This page may be photocopied noncommercially by physicians and other health care professionals to share with patients. To purchase bulk reprints, call 203/259-8724.

FOR MORE INFORMATION

- National Institute of Diabetes and Digestive and Kidney Diseases
www.niddk.nih.gov
- American Liver Foundation
800/465-4837
www.liverfoundation.org
- United Network for Organ Sharing
888/894-6361
www.unos.org

INFORM YOURSELF

To find this and previous JAMA Patient Pages, go to the Patient Page link on JAMA's Web site at www.jama.com. Many are available in English and Spanish. A Patient Page on hepatitis C was published in the May 14, 2003, issue; and one on suppressing the immune system for organ transplants was published in the May 10, 2000, issue.

