



Pulmonary Embolism

Sharon Parmet; Cassio Lynn; Richard M. Glass

Online article and related content
current as of July 5, 2009.

JAMA. 2006;295(2):240 (doi:10.1001/jama.295.2.240)

<http://jama.ama-assn.org/cgi/content/full/295/2/240>

Supplementary material

Spanish PDF

<http://jama.ama-assn.org/cgi/content/full/295/2/240/DC1>

Correction

[Contact me if this article is corrected.](#)

Citations

[Contact me when this article is cited.](#)

Topic collections

Venous Thromboembolism; Cardiovascular System; JAMA Patient Page
[Contact me when new articles are published in these topic areas.](#)

Related Articles published in
the same issue

Effectiveness of Managing Suspected Pulmonary Embolism Using an
Algorithm Combining Clinical Probability, D-Dimer Testing, and
Computed Tomography
Writing Group for the Christopher Study Investigators. JAMA. 2006;295(2):172.

Subscribe

<http://jama.com/subscribe>

Email Alerts

<http://jamaarchives.com/alerts>

Permissions

permissions@ama-assn.org

<http://pubs.ama-assn.org/misc/permissions.dtl>

Reprints/E-prints

reprints@ama-assn.org

Pulmonary Embolism

A **pulmonary embolism** is a blood clot that travels to the lungs. Often, the clot forms in another part of the body, usually in the veins of the legs. Large clots can be fatal. The January 11, 2006, issue of *JAMA* includes an article about diagnosing pulmonary embolism. This Patient Page is based on one previously published in the December 3, 2003, issue of *JAMA*.

SYMPTOMS AND SIGNS OF PULMONARY EMBOLISM

- Sudden chest pain
- Difficulty breathing
- Cough with blood
- Lightheadedness and fainting
- Excessive sweating
- Bluish tint to skin
- Anxiety

RISK FACTORS FOR PULMONARY EMBOLISM

- Recent surgery
- Long periods of immobility (such as bed rest because of an illness)
- Cancer
- Heart problems
- Older age
- Taking oral contraceptives
- Previous stroke or heart attack

DIAGNOSIS OF PULMONARY EMBOLISM

Pulmonary embolism can be difficult to diagnose because the symptoms are broad and resemble those of other diseases. Results of the patient's history and physical examination are very important. Some other tests that may be done include

- Chest x-ray
- Electrocardiogram
- Arterial blood gases—measurement of oxygen and carbon dioxide levels in the blood
- D-dimer assay—a blood test for evidence of blood clots
- Ultrasound of the legs—sound wave images to detect blood clots in the veins
- Spiral computed tomography—detailed computerized x-ray imaging
- Lung scan—test of blood flow through the lungs
- Pulmonary arteriogram—injections to show the arteries in the lungs to detect blood clots

TREATMENT

Immediate treatment usually includes giving injectable **anticoagulants** (blood thinners)—such as **heparin**, followed by **warfarin** (an anticoagulant taken by mouth)—to stabilize the clot, prevent additional clots, and restore normal blood flow in the lungs. Oxygen and sedatives may be given to make the patient more comfortable. If the clot is large, **thrombolytic** ("clot busting") drugs may be needed to remove it.

Sources: American Heart Association; National Heart, Lung, and Blood Institute

Sharon Parmet, MS, Writer

Cassio Lynn, MA, Illustrator

Richard M. Glass, MD, Editor

The JAMA Patient Page is a public service of JAMA. The information and recommendations appearing on this page are appropriate in most instances, but they are not a substitute for medical diagnosis. For specific information concerning your personal medical condition, JAMA suggests that you consult your physician. This page may be photocopied noncommercially by physicians and other health care professionals to share with patients. To purchase bulk reprints, call 718/946-7424.

FOR MORE INFORMATION

- American Heart Association
www.americanheart.org
- National Heart, Lung, and Blood Institute
www.nhlbi.nih.gov

INFORM YOURSELF

To find this and other JAMA Patient Pages, go to the Patient Page link on JAMA's Web site at www.jama.com. A previous Patient Page on thrombophlebitis was published in the August 10, 2005, issue.

