



Coronary Artery Bypass Grafting

Sharon Parmet; Cassio Lynn; Richard M. Glass

JAMA. 2008;299(15):1856 (doi:10.1001/jama.299.15.1856)

Online article and related content
current as of November 9, 2009.

<http://jama.ama-assn.org/cgi/content/full/299/15/1856>

Supplementary material

Spanish PDF

<http://jama.ama-assn.org/cgi/content/full/299/15/1856/DC1>

Correction

[Contact me if this article is corrected.](#)

Citations

[Contact me when this article is cited.](#)

Topic collections

Revascularization; Cardiovascular System; Surgery; Surgical Interventions; Cardiovascular/ Cardiothoracic Surgery; JAMA Patient Page; Cardiovascular Intervention

[Contact me when new articles are published in these topic areas.](#)

Subscribe

<http://jama.com/subscribe>

Email Alerts

<http://jamaarchives.com/alerts>

Permissions

permissions@ama-assn.org

<http://pubs.ama-assn.org/misc/permissions.dtl>

Reprints/E-prints

reprints@ama-assn.org

Coronary Artery Bypass Grafting

Coronary artery bypass grafting (CABG) is a surgical procedure that uses another artery or vein to reroute blood around a blockage in the arteries (**coronary arteries**) that supply the heart with blood and oxygen. The April 16, 2008, issue of *JAMA* includes an article about CABG surgery. This Patient Page is adapted from one published in the April 21, 2004, issue of *JAMA*.

CORONARY ARTERY DISEASE

- In persons with coronary artery disease (CAD), deposits of cholesterol and fats called **plaque** form in the coronary arteries. This process is called **atherosclerosis** (hardening of the arteries).
- If plaque continues to build up, blood vessels can become partially or completely blocked so the heart does not receive enough oxygen carried by red blood cells, leading to **angina** (chest pain) or even a **myocardial infarction** (heart attack).
- CABG surgery may be done to bypass blocked coronary arteries.

CABG SURGERY PROCEDURES

- Before undergoing CABG surgery, a special dye that can be seen on x-ray film is injected into the coronary arteries while x-rays are taken.
- Blood-thinning drugs called **anticoagulants** are given to help prevent blood clot formation during the operation.
- If a heart-lung bypass machine is used, the machine takes over pumping blood for the heart (conventional "on-pump" CABG).
- If a heart-lung bypass machine is not used, the heart is positioned and the coronary artery to be bypassed is stabilized with special devices ("beating heart" or "off-pump" CABG).
- Another artery or vein is sewn into place so that blood flow can bypass the blockage in the coronary artery. If more than one artery is blocked, more blood vessels will be used to bypass them (called **double**, **triple**, or **quadruple bypass surgery**).

RECOVERING FROM SURGERY

- After CABG surgery, the patient recovers in the intensive care unit (ICU).
- Many patients can go home about 3 to 6 days after surgery.
- It will take another 4 to 6 weeks for the patient to feel stronger and resume his or her normal activities.


FOR MORE INFORMATION

- American Heart Association
www.americanheart.org
- Cardiothoracic Surgery Network
www.ctsnet.org

INFORM YOURSELF

To find this and other JAMA Patient Pages, go to the Patient Page link on JAMA's Web site at www.jama.com. A Patient Page on acute coronary syndromes was published in the August 15, 2007, issue.

Sources: American Heart Association; National Heart, Lung, and Blood Institute; Heart Information Network



Sharon Parmet, MS, Writer

Cassio Lynn, MA, Illustrator

Richard M. Glass, MD, Editor

The JAMA Patient Page is a public service of JAMA. The information and recommendations appearing on this page are appropriate in most instances, but they are not a substitute for medical diagnosis. For specific information concerning your personal medical condition, JAMA suggests that you consult your physician. This page may be photocopied noncommercially by physicians and other health care professionals to share with patients. To purchase bulk reprints, call 203/259-8724.

