



Weight and Diabetes

Sharon Parmet; Cassio Lynn; Richard M. Glass

JAMA. 2008;299(23):2814 (doi:10.1001/jama.299.23.2814)

<http://jama.ama-assn.org/cgi/content/full/299/23/2814>

Online article and related content
current as of November 22, 2009.

Supplementary material

Spanish PDF

<http://jama.ama-assn.org/cgi/content/full/299/23/2814/DC1>

Correction

[Contact me if this article is corrected.](#)

Citations

[Contact me when this article is cited.](#)

Topic collections

Public Health; Obesity; JAMA Patient Page; Endocrine Diseases;
Diabetes Mellitus

[Contact me when new articles are published in these topic areas.](#)

Related Articles published in
the same issue

Examining a Bidirectional Association Between Depressive Symptoms
and Diabetes

Sherita Hill Golden et al. *JAMA*. 2008;299(23):2751.

Subscribe

<http://jama.com/subscribe>

Email Alerts

<http://jamaarchives.com/alerts>

Permissions

permissions@ama-assn.org

<http://pubs.ama-assn.org/misc/permissions.dtl>

Reprints/E-prints

reprints@ama-assn.org

Weight and Diabetes

Diabetes is a common disorder in which the body has difficulty controlling levels of sugar in the bloodstream. Normally, the hormone **insulin** made by the **pancreas** (an organ in the abdomen) regulates blood sugar levels. The bodies of individuals with **type 1 diabetes**, which usually starts by the early teen years, do not make enough insulin to control blood sugar, so they must receive insulin injections. The bodies of persons with type 2 diabetes are resistant to the effects of insulin. **Type 2 diabetes**, also known as “adult-onset” diabetes, usually develops in adulthood but can also occur in overweight children. Family history of diabetes and excess weight, especially weight carried around the middle, are strong risk factors for developing type 2 diabetes. Losing weight greatly reduces your chances for type 2 diabetes and can help bring your blood sugar under control if you already have type 2 diabetes. Type 2 diabetes can be treated with diet, exercise, and oral prescription medications but may require insulin shots.

The June 18, 2008, issue of JAMA includes an article about an association between type 2 diabetes and depression. This Patient Page is based on one previously published in the March 15, 2006, issue of JAMA.



TO LOSE OR CONTROL WEIGHT

- Get regular exercise—at least 30 minutes per day most days of the week of brisk walking, sports, or active games.
- Eat a healthful diet rich in vegetables, fruits, and whole grains and low in refined carbohydrates, such as sweets and white bread.
- Limit the amount of high-sugar beverages you drink, such as soft drinks and fruit punches.
- Avoid high-fat foods like ice cream, butter, and high-fat meats.
- Limit alcohol to no more than 1 drink per day for women, 2 per day for men, and none if you have any difficulty controlling alcohol intake.
- Always eat a balanced breakfast.
- If you are overweight, aim to lose no more than 2 pounds per week—losing more than that can be unhealthy and often leads to rebound weight gain.
- Get your family and friends involved by encouraging them to eat healthful foods and exercise together.

Realize that your diet and exercise regimen are lifestyle changes that must be maintained in the long term to keep weight off.

Sources: National Institute of Diabetes and Digestive and Kidney Diseases, American Diabetes Association, Centers for Disease Control and Prevention

Sharon Parmet, MS, Writer

Cassio Lynn, MA, Illustrator

Richard M. Glass, MD, Editor

The JAMA Patient Page is a public service of JAMA. The information and recommendations appearing on this page are appropriate in most instances, but they are not a substitute for medical diagnosis. For specific information concerning your personal medical condition, JAMA suggests that you consult your physician. This page may be photocopied noncommercially by physicians and other health care professionals to share with patients. To purchase bulk reprints, call 312/464-0776.

FOR MORE INFORMATION

- American Diabetes Association
www.diabetes.org
- National Institute of Diabetes and Digestive and Kidney Diseases
www.niddk.nih.gov

INFORM YOURSELF

To find this and other JAMA Patient Pages, go to the Patient Page link on JAMA's Web site at www.jama.com. A Patient Page on depression was published in the May 28, 2008, issue; one on type 2 diabetes in children was published in the September 26, 2001, issue; one on the ABCs of diabetes was published in the May 15, 2002, issue; and one on type 1 diabetes was published in the October 22/29, 2003, issue.

JAMA
COPY FOR
YOUR PATIENTS