



Thrombophlebitis

Janet M. Torpy; Alison E. Burke; Richard M. Glass

Online article and related content
current as of November 23, 2009.

JAMA. 2008;300(14):1718 (doi:10.1001/jama.300.14.1718)

<http://jama.ama-assn.org/cgi/content/full/300/14/1718>

Supplementary material

Spanish PDF

<http://jama.ama-assn.org/cgi/content/full/300/14/1718/DC1>

Correction

[Contact me if this article is corrected.](#)

Citations

[Contact me when this article is cited.](#)

Topic collections

Venous Thromboembolism; Cardiovascular System; JAMA Patient Page

[Contact me when new articles are published in these topic areas.](#)

Related Articles published in
the same issue

Serial 2-Point Ultrasonography Plus D-Dimer vs Whole-Leg Color-Coded
Doppler Ultrasonography for Diagnosing Suspected Symptomatic Deep
Vein Thrombosis: A Randomized Controlled Trial

Enrico Bernardi et al. *JAMA*. 2008;300(14):1653.

Subscribe

<http://jama.com/subscribe>

Email Alerts

<http://jamaarchives.com/alerts>

Permissions

permissions@ama-assn.org

<http://pubs.ama-assn.org/misc/permissions.dtl>


Reprints/E-prints

reprints@ama-assn.org

Thrombophlebitis

Blood clots can form in arteries or veins. When inflammation due to a blood clot occurs in a vein, it is called **thrombophlebitis**.

Thrombophlebitis usually occurs in the veins of the legs. Rarely, it can occur in the veins of the arms or neck. **Superficial** (near the surface) thrombophlebitis occurs in the veins that are visible just under the skin. The area of inflammation is usually reddened, tender, and warm to the touch and can be painful. The leg or arm may swell and fever may occur. **Deep venous thrombosis (DVT)** is more dangerous than superficial thrombophlebitis. Deep venous thrombosis often cannot be seen or felt by the individual. Swelling of the leg or arm or fever may alert a person to the presence of a DVT, especially if risk factors for DVT exist. **Pulmonary embolism**, a condition that can be fatal, results from a DVT that becomes loose in the venous system and travels to the lungs. There, the blood clot blocks proper blood flow to the lungs and decreases oxygen levels in the body. The October 8, 2008, issue of *JAMA* includes an article about diagnostic testing for DVT. This Patient Page is based on one previously published in the July 26, 2006, issue of *JAMA*.



RISK FACTORS

- Inactivity due to recent injury, surgery, or prolonged sitting
- Pregnancy or recent childbirth
- Oral contraceptive use or estrogen therapy
- Cancer
- Stroke or other diseases that limit movement
- Family history of clotting disorders
- Central venous catheters (used for injection of medications or for dialysis)

SYMPTOMS

In addition to a medical history and physical examination, the doctor may order tests to evaluate superficial thrombophlebitis or to look for presence of a DVT. These tests may include **ultrasound** (using sound waves to look for a blood clot in the vein), computed tomography (CT) or magnetic resonance imaging (MRI) to visualize blood vessels, blood tests, or **venography** (using injection of a dye to trace the course of a vein).

PREVENTION

- Move your legs, especially during prolonged sitting or bed rest
- Use compression (strong support) stockings
- Discuss your personal and family history with your doctor before considering hormone therapy

TREATMENT

Treatment for superficial thrombophlebitis usually includes elevating the leg, warm compresses to the area, and medication to decrease pain and inflammation. Support stockings may be worn to reduce swelling. Treatment for DVT or pulmonary embolism usually involves **anticoagulation** (blood thinning) treatment with **heparin** (by injection) or **warfarin** (by mouth for longer-term treatment). Pregnant women should not use warfarin because it can harm the developing fetus.

FOR MORE INFORMATION

- National Heart, Lung, and Blood Institute
www.nhlbi.nih.gov
- American Heart Association
www.americanheart.org

INFORM YOURSELF

To find this and previous JAMA Patient Pages, go to the Patient Page Index on JAMA's Web site at www.jama.com. Many are available in English, Spanish, and French. Patient Pages on pulmonary embolism were published in the December 3, 2003, and the February 14, 2001, issues.

Sources: National Heart, Lung, and Blood Institute; American Heart Association

Janet M. Torpy, MD, Writer

Alison E. Burke, MA, Illustrator

Richard M. Glass, MD, Editor

The JAMA Patient Page is a public service of JAMA. The information and recommendations appearing on this page are appropriate in most instances, but they are not a substitute for medical diagnosis. For specific information concerning your personal medical condition, JAMA suggests that you consult your physician. This page may be photocopied noncommercially by physicians and other health care professionals to share with patients. To purchase bulk reprints, call 312/464-0776.

