

Binge Drinking Among US Adults

Timothy S. Naimi, MD, MPH

Robert D. Brewer, MD, MSPH

Ali Mokdad, PhD

Clark Denny, PhD

Mary K. Serdula, MD, MPH

James S. Marks, MD, MPH

ALCOHOL ABUSE, WHICH KILLS approximately 100 000 Americans annually, is the third leading preventable cause of death in the United States.¹ *Binge drinking*, generally defined as the consumption of 5 or more alcoholic beverages on 1 occasion,^{2,3} generally results in acute impairment and causes a substantial fraction of all alcohol-related deaths.^{4,5} Adverse health effects specifically associated with binge drinking include unintentional injuries (eg, motor vehicle crashes, falls, drowning, hypothermia, and burns), suicide, sudden infant death syndrome⁶ alcohol poisoning, hypertension, acute myocardial infarction, gastritis, pancreatitis, sexually transmitted diseases, meningitis, and poor control of diabetes. Many consequences of binge drinking have especially high social and economic costs, including interpersonal violence (eg, homicide, assault, domestic violence, rape, and child abuse), fetal alcohol syndrome, unintended pregnancy, child neglect, and lost productivity.^{7,8}

Reducing binge drinking among adults is one of the leading health indicators in Healthy People 2010.⁹ Recent reports have indicated that prior declines in alcohol consumption reached a plateau in the early 1990s (M. K. S., written communication, December 12, 2002)¹⁰ and that the preva-

Context Binge drinking (consuming ≥ 5 alcoholic drinks on 1 occasion) generally results in acute impairment and has numerous adverse health consequences. Reports indicate that binge drinking may be increasing in the United States.

Objectives To quantify episodes of binge drinking among US adults in 1993-2001, to characterize adults who engage in binge drinking, and to describe state and regional differences in binge drinking.

Design, Setting, and Participants The Behavioral Risk Factor Surveillance System, a random-digit telephone survey of adults aged 18 years or older that is conducted annually in all states. The sample size ranged from 102 263 in 1993 to 212 510 in 2001.

Main Outcome Measures Binge-drinking prevalence, episodes, and episodes per person per year.

Results Between 1993 and 2001, the total number of binge-drinking episodes among US adults increased from approximately 1.2 billion to 1.5 billion; during this time, binge-drinking episodes per person per year increased by 17% (from 6.3 to 7.4, P for trend = .03). Between 1995 and 2001, binge-drinking episodes per person per year increased by 35% (P for trend = .005). Men accounted for 81% of binge-drinking episodes in the study years. Although rates of binge-drinking episodes were highest among those aged 18 to 25 years, 69% of binge-drinking episodes during the study period occurred among those aged 26 years or older. Overall, 47% of binge-drinking episodes occurred among otherwise moderate (ie, non-heavy) drinkers, and 73% of all binge drinkers were moderate drinkers. Binge drinkers were 14 times more likely to drive while impaired by alcohol compared with non-binge drinkers. There were substantial state and regional differences in per capita binge-drinking episodes.

Conclusions Binge drinking is common among most strata of US adults, including among those aged 26 years or older. Per capita binge-drinking episodes have increased, particularly since 1995. Binge drinking is strongly associated with alcohol-impaired driving. Effective interventions to prevent the mortality and morbidity associated with binge drinking should be widely adopted, including screening patients for alcohol abuse in accordance with national guidelines.

JAMA. 2003;289:70-75

www.jama.com

lence of binge drinking and adverse health outcomes associated with binge drinking have increased recently.^{11,12} Because people who binge drink may do so repeatedly, the percentage of adults who binge drink represents only the tip of the iceberg relative to the actual number of binge-drinking episodes among US adults. We used data from the Behavioral Risk Factor Surveillance System (BRFSS) survey from 1993 to 2001 to quantify episodes of binge drinking among US adults and to characterize adult binge drinkers. Although BRFSS

has been used to estimate the prevalence of binge drinking, we are not aware of any previous studies that re-

Author Affiliations: From the Alcohol Team, Emerging Investigations, and Analytic Methods Branch (Drs Naimi and Brewer); Division of Applied Public Health Training, Epidemiology Program Office (Dr Naimi); and Behavioral Surveillance Branch, Division of Adult and Community Health (Drs Mokdad and Denny); Chronic Disease Nutrition Branch, Division of Nutrition and Physical Activity (Dr Serdula); Office of the Director, National Center for Chronic Disease Prevention and Health Promotion (Dr Marks); Centers for Disease Control and Prevention, Atlanta, Ga.

Corresponding Author and Reprints: Timothy S. Naimi, CDC/Alcohol Team, MS K-67, 4770 Buford Hwy, NE, Atlanta, GA 30341 (e-mail: tbn7@cdc.gov).

port population-based estimates of binge-drinking episodes in the US. Furthermore, the large sample size of BRFSS provided a unique opportunity to generate state-specific estimate of per capita binge-drinking episodes and enabled us to assess binge drinkers by a variety of demographic and sociocultural factors.

METHODS

The BRFSS, a series of state-based health surveys coordinated by the Centers for Disease Control and Prevention, in conjunction with state health departments, is a cross-sectional random-digit telephone survey of US adults 18 years or older who are not institutionalized. Details of the sampling, purpose, and method of analysis have been published elsewhere.¹³⁻¹⁵ We analyzed BRFSS data from 49 states for 1993 and from 50 states for 1995, 1997, 1999, and 2001. Wyoming, the 50th BRFSS state, which was added in 1994, accounted for less than 0.2% of binge-drinking episodes in those years in which it was included.

The alcohol section of the BRFSS questionnaire first asks respondents whether they had any alcoholic beverages in the past 30 days. Those who respond yes are asked how frequently they drank alcohol (measured as days per week or month) and how many drinks they had on days they drank. Separately, they also are asked how often they consume 5 or more drinks on a single occasion (ie, binge-drinking episodes) in the past 30 days. The BRFSS survey itself does not use the term *binge drinking*.

Binge drinking was defined as the consumption of 5 or more drinks on at least 1 occasion in the previous 30 days.³ We calculated binge-drinking episodes per respondent per year by multiplying the number of self-reported binge-drinking episodes for each respondent by 12. Because BRFSS interviews a representative, stratified sample of US adults each month, combining all data for 12 months yields a representative sample for the entire year. In addition, seasonal variations in

binge drinking are accounted for when episodes reported for 12 months are aggregated for all respondents. To generate estimates of binge-drinking episodes per capita (ie, episodes per person per year) for various population strata, we divided the number of binge-drinking episodes for each stratum by census-based estimates of the relevant population denominator for the year in question.

Moderate drinking was defined as consuming an average of 2 drinks per day or less for a man and 1 drink per day or less for a woman.¹⁶ *Heavy drinking* was defined as consuming alcohol in excess of moderate levels. To calculate average daily drinks, we multiplied the proportion of days the respondent drank alcohol by the average number of drinks on days when alcohol was consumed, and divided by 30. Alcohol-impaired driving was defined as a non-zero response to the question: "During the past 30 days, how many times have you driven after having had perhaps too much [alcohol] to drink?"

The BRFSS sample size ranged from 102 263 for 1993 to 212 510 for 2001, with those not providing information on alcohol consumption excluded from analysis. For the study period, the response rate (completed interviews divided by the sum of completed interviews, terminated interviews, and refusals) ranged from 68.4% to 82.1%. Analyses were conducted using SAS (SAS Institute Inc, Cary, NC) and SUDAAN software.¹⁷

RESULTS

Between 1993 and 2001, the total number of binge-drinking episodes among US adults increased from approximately 1.2 billion to 1.5 billion; during this time, binge-drinking episodes per person per year increased by 17% (from 6.3 to 7.4, P for trend = .03) (TABLE 1 and FIGURE 1). Between 1995 and 2001, binge-drinking episodes per person per year increased by 35% (P for trend = .005). During the study period, similar changes were noted for most population strata by sex, age, race/ethnicity, and education. In 2001,

14.3% of US adults had at least 1 binge-drinking episode during the previous 30 days.

Binge drinking rates were highest among younger age groups, and declined with increasing age (Table 1). In 2001, young adults aged 21 to 25 years (18.0 episodes/person/year) and underage drinkers aged 18 to 20 years (15.3 episodes/person/year) had the highest rates of binge-drinking episodes. During the study period, underage drinkers had the largest increase (56%) in the rate of binge-drinking episodes. However, the rate of binge-drinking episodes among respondents aged 26 to 55 years also increased by approximately 25% during this period. Those older than 55 years had the lowest rate of binge-drinking episodes (<3 episodes/person/year). Although older age groups had lower rates of binge drinking, the majority of binge-drinking episodes occurred among those who were 26 years or older (Figure 1), with 69% of binge-drinking episodes occurring among this same age group.

During the study period, men accounted for 81% of all binge-drinking episodes (Table 1), and they were approximately 3 times as likely as women to report binge drinking (eg, 22.7% vs 7.2% in 2001). However, among women of childbearing age (18-44), 11.8% reported binge drinking; each woman who binge drank averaged 36.6 episodes in 2001 (results not shown in Table 1). Whites accounted for 78% of all binge-drinking episodes (Table 1), although Hispanics had the highest rate of binge-drinking episodes per person for most years of the study (eg, 8.4 episodes/person/year in 2001). Blacks consistently had the lowest rate of binge-drinking episodes per capita, averaging less than 5 episodes per person per year. College graduates had slightly lower rates of binge drinking compared with other educational strata. Heavy drinkers were more likely to binge drink and had far higher rates of binge-drinking episodes than moderate drinkers. However, moderate drinkers accounted for 47% of binge-drinking episodes, including 639 million binge-drinking episodes in 2001.

Table 1. Prevalence, Total Number, and Rate of Binge-Drinking Episodes Among US Adults, 1993-2001*


Characteristics	Binge Drinkers, %†					No. Binge-Drinking Episodes (Millions/Year)					Rate of Binge-Drinking Episodes (Episodes/Person/Year)				
	1993	1995	1997	1999	2001	1993	1995	1997	1999	2001	1993	1995	1997	1999	2001
	(n = 102 464)	(n = 113 934)	(n = 135 582)	(n = 159 989)	(n = 212 510)										
All respondents	14.2	12.7	13.2	14.6	14.3	1179	945	1170	1334	1520	6.3	5.5	6.1	6.9	7.4
Sex															
Male	21.8	19.9	20.8	22.7	22.2	961	765	955	1078	1235	10.8	9.40	10.5	11.6	12.5
Female	7.3	6.3	6.3	7.1	7.2	218	180	215	256	285	2.2	2.0	2.2	2.5	2.7
Age, y															
18-20	23.0	19.6	22.0	26.0	26.1	94	76	116	132	157	9.8	8.8	12.0	13.7	15.3
21-25	30.0	26.5	27.4	32.2	32.2	279	196	236	276	339	14.1	11.6	13.4	15.6	18.0
26-34	20.7	19.5	20.2	21.3	21.0	320	248	275	299	314	8.4	7.6	7.9	9.0	9.2
35-54	12.4	11.7	12.5	13.6	13.6	355	327	419	468	539	5.4	5.1	5.7	6.2	6.7
≥55	4.2	3.8	3.9	4.7	4.3	130	95	122	158	166	2.5	1.9	2.1	2.8	2.7
Race/ethnicity															
White	14.7	13.4	13.6	14.9	14.7	971	778	918	1026	1098	6.6	5.8	6.4	7.1	7.4
Black	8.9	8.7	9.0	10.2	9.8	69	65	74	89	106	4.2	4.0	4.0	4.8	5.4
Hispanic	16.1	11.7	15.8	17.2	16.8	105	69	120	160	197	7.0	5.1	6.4	7.2	8.4
Other	11.3	10.1	10.8	12.2	13.4	31	25	51	48	108	5.2	4.1	6.0	6.3	8.2
Education															
Some high school	10.7	8.5	10.3	11.8	12.8	162	123	143	181	209	6.0	4.7	5.5	6.8	8.0
High school graduate	14.3	13.3	13.2	14.8	14.4	412	331	409	481	522	6.6	5.8	6.6	7.7	8.2
Some college	16.2	15.0	15.2	16.4	16.2	357	291	380	389	462	7.3	6.4	7.2	7.5	8.3
College graduate	13.9	12.2	12.8	13.8	13.3	247	199	237	282	325	5.2	4.6	4.7	5.4	5.4
Alcohol intake‡															
Moderate	23.1	23.0	21.6	23.1	21.6	587	485	539	608	639	6.7	6.5	6.3	6.9	6.4
Heavy	73.7	74.8	67.2	68.4	76.0	592	459	630	725	881	70.1	69.8	60.7	62.2	81.9

*Binge drinking is defined as consuming ≥5 alcohol-containing drinks on 1 occasion.

†Percentage of Behavioral Risk Factor Surveillance System respondents who reported binge drinking in the past 30 days.

‡Moderate alcohol intake is defined as consuming an average of ≤1 alcohol-containing drink per day for a woman or ≤2 for a man; heavy alcohol intake, consuming an average of >1 alcohol-containing drink per day for a woman or >2 for a man.

Figure 1. Binge-Drinking Episodes by Age in US Adults, 1993-2001



In 2001, 101 million (49%) of 205 million American adults consumed alcohol (data not shown). More than 60% of adult male drinkers 18 to 25 years

reported binge drinking; males in this age group averaged 39 binge-drinking episodes in 2001 (TABLE 2). By race/ethnicity, Hispanic drinkers had the highest prevalence of binge drinking, including 45.3% of Hispanic males. Overall, 22% of moderate-drinking US adults reported binge drinking, including 30.1% of male adults who were moderate drinkers. In 2001, all US moderate drinkers averaged 6.4 binge-drinking episodes per capita. Among heavy drinkers, 76% reported binge drinking, and each heavy drinker averaged approximately 1.5 binge-drinking episodes each week.

Among US adults who consumed alcohol in 2001, binge drinkers were younger, more likely to be male, and less likely to have had any college education compared with non-binge drinkers (TABLE 3). Although binge drinkers consumed an average of approximately

1 drink per day more than non-binge drinkers, 72.9% of all binge drinkers were moderate drinkers. Binge drinkers also were approximately 14 times as likely to report alcohol-impaired driving than non-binge drinkers in 1999 (this question was not asked in the BRFSS survey in 2001).

In 2001, rates of binge-drinking episodes varied considerably by state. FIGURE 2 shows unadjusted rates for all states, which ranged from 3.1 episodes per person per year in Tennessee to 12.3 episodes per person per year in Wisconsin. Overall, per capita rates were higher in the northcentral and western US, and lower in the southeastern US. To determine the extent to which state and regional differences were due to demographic factors, we adjusted the age, sex, and race/ethnicity data for each state to the US population. After adjustment, the range of episodes per person per year

(3.9-12.5) was similar to that of the unadjusted data, and the state and regional differences persisted. Using the cut points chosen to generate Figure 2, 39 (78%) of 50 states remained in the same category as that determined by the unadjusted data. Adjusted rates changed 3 states (Florida, New York, and Texas) from the intermediate category of binge drinking to the highest category, changed 1 state (Iowa) from highest to intermediate, changed 4 states (Arkansas, Louisiana, Maryland, and Washington) from lowest to intermediate, changed 2 states (New Mexico and Colorado) from intermediate to lowest, and changed 2 states (Montana and Wyoming) from highest to lowest.

COMMENT

To our knowledge, this is the first US study to quantify binge-drinking episodes. Overall, there were approximately 1 billion or more episodes of binge drinking annually from 1993 to 2001. Especially high rates of binge-drinking episodes were observed among men, young adults, and heavy drinkers. However, about 70% of binge-drinking episodes were reported among those 26 years or older, and approximately half of all binge-drinking episodes were reported among moderate (ie, non-heavy) drinkers.

Our finding that binge drinking rates are highest among young age groups is consistent with other studies.^{10,18} However, we found that binge drinking is by no means confined to young age groups. Although those aged 18 to 25 years had the highest number of binge-drinking episodes per capita, rates remained high throughout the study period (approximately 8 episodes per year) among respondents aged 26 to 34 years, contributing to the large proportion (70%) of total episodes that were observed among persons 26 years or older. Although efforts to prevent youth access to alcohol and binge drinking on college campuses are critical, they need to be complemented by interventions that will reduce binge drinking among adults of all ages, particularly those younger than 55 years.

Our finding that there are almost as many binge drinking episodes among moderate drinkers as heavy drinkers has been noted in other countries,¹⁹ and is largely because there are more moderate drinkers than heavy drinkers in the general population. However, these findings also emphasize why it is important to assess binge alcohol use independent of average daily alcohol con-

sumption, and underscores why binge drinking is a key indicator of alcohol abuse among both moderate and heavy drinkers.^{7,20}

Although males accounted for most binge-drinking episodes, the high rate of binge drinking reported among women of childbearing age is concerning because of alcohol's harmful effects on the developing fetus, particu-

Table 2. Binge Drinking Among US Adults Who Consumed Alcohol, 2001*

Characteristic	Males (n = 57 654)		Females (n = 46 811)		Total (N = 104 465)	
	%†	Rate‡	%†	Rate‡	%	Rate
All respondents	35.9	20.1	15.7	5.8	26.8	13.7
Age, y						
18-20	61.1	39.0	37.7	17.6	51.3	30.0
21-25	61.9	38.7	32.0	12.5	48.6	27.1
26-34	44.2	20.8	20.2	6.5	34.1	14.8
35-54	33.3	17.9	13.5	4.7	24.2	11.9
≥55	15.0	10.4	4.7	1.8	10.2	6.4
Race/ethnicity						
White	34.6	19.3	15.6	5.6	25.8	13.0
Black	33.4	20.7	14.1	5.1	24.4	13.4
Hispanic	45.3	23.5	18.1	7.0	35.1	17.3
Other	37.2	24.1	15.3	6.6	29.0	17.6
Education						
Some high school	45.3	29.8	23.1	10.1	37.5	23.0
High school graduate	41.9	26.3	16.6	6.2	30.7	17.4
Some college	38.5	21.6	17.8	7.0	28.3	14.4
College graduate	27.0	11.9	11.7	3.6	20.3	8.2
Alcohol intake§						
Moderate	30.1	9.5	11.3	2.5	21.6	6.4
Heavy	88.2	113.8	59.6	37.9	76.0	81.9

*Binge drinking is defined as consuming ≥5 alcohol-containing drinks on 1 occasion.

†Percentage of US adults drinkers who had at least 1 binge-drinking episode in the past 30 days.

‡Number of episodes of binge drinking per person per year (among drinkers for the given demographic group).

§Moderate alcohol intake is defined as consuming an average of ≤1 alcohol-containing drink per day for a woman or ≤2 for a man; heavy alcohol intake, consuming an average of >1 alcohol-containing drink per day for a woman or >2 for a man.

Table 3. US Adults Who Consumed Alcohol, by Binge-Drinking Status, 2001

Characteristic	Binge Drinkers* (n = 27 997)	Non-Binge Drinkers (n = 76 468)	Total (N = 104 465)
Age, mean (SE), y	35.1 (0.15)	45.5 (0.10)	42.5 (0.09)
Male, %	73.8	48.4	55.1
White, %	74.1	78.1	77.1
College, %†	57.8	67.6	65.0
Moderate drinker, %‡	72.9	96.9	90.4
Mean (SE) drinks/day, No.	1.6 (0.02)	0.4 (0.005)	0.7 (0.01)
Impaired driver, %§	13.8	1.0	4.6

*Binge drinking is defined as consuming ≥5 alcohol-containing drinks on 1 occasion.

†At least some college education.

‡Moderate alcohol intake is defined as consuming an average of ≤1 alcohol-containing drink per day for a woman or ≤2 for a man; heavy alcohol intake, consuming an average of >1 alcohol-containing drink per day for a woman or >2 for a man.

§Impaired drivers were those drinkers who provided a non-zero response to the question: "During the past 30 days, how many times have you driven after having had perhaps too much [alcohol] to drink?" Impaired driving estimates are from 1999 data, since the impaired driving questions were not asked in 2001.

- G, Chalmers N. Quantifying the disease impact of alcohol with ARDI software. *Public Health Rep.* 1991; 106:443-450.
9. US Department of Health and Human Services. *Healthy People 2010*. Vol 1. 2nd ed. Washington, DC: US Government Printing Office; 2000.
10. Greenfield TK, Midanik LT, Rogers JD. A 10-year national trend study of alcohol consumption, 1984-1995: is the period of declining drinking over? *Am J Public Health.* 2000;90:47-52.
11. Nelson DE, Bland S, Powell-Griner E, et al. State trends in health risk factors and receipt of clinical preventive services among US adults during the 1990s. *JAMA.* 2002;287:2659-2667.
12. Centers for Disease Control and Prevention. Alcohol involvement in fatal motor-vehicle crashes—United States, 1990-2000. *MMWR Morb Mortal Wkly Rep.* 2001;50:1064-1065.
13. Gentry EM, Kalsbeek WD, Hogelin GC, et al. The behavioral risk factor surveys, II. *Am J Prev Med.* 1985; 1:9-14.
14. Remington PL, Smith MY, Williamson DF, Anda RF, Gentry EM, Hogelin G. Design, characteristics, and usefulness of state-based behavioral risk factor surveillance: 1981-1987. *Public Health Rep.* 1988;103: 366-375.
15. Waksberg JS. Methods for random digit dialing. *J Am Stat Assoc.* 1978;73:40-46.
16. US Department of Health and Human Services, US Department of Agriculture. *Nutrition and Your Health: Dietary Guidelines for Americans*. 5th ed. Home and Gardening Bulletin, No. 232. Washington, DC: US Government Printing Office; 2000.
17. Shah BV, Bieler GS. *SUDAAN User's Manual, release 7.5*. Research Triangle Park, NC: Research Triangle Institute; 1997.
18. Substance Abuse and Mental Health Services Administration. 1999 National Household on Drug Abuse summary findings. Available at: <http://www.samhsa.gov/oas/nhsda.htm>. Accessed November 26, 2002.
19. Gmel G, Klingemann S, Muller R, Brenner D. Revisiting the preventive paradox: the Swiss case. *Addiction.* 2001;96:273-284.
20. World Health Organization. *International Guide for Monitoring Alcohol Consumption and Related Harms*. Geneva, Switzerland: World Health Organization; 2000.
21. Jacobson JL, Jacobson SW. Prenatal alcohol exposure and neurobehavioral development: where is the threshold? *Alcohol Health Res World.* 1994;18:30-36.
22. Maier SE, Chen WA, Miller JA, West JR. Fetal alcohol exposure and temporal vulnerability: regional differences in alcohol-induced microencephaly as a function of the timing of binge-like alcohol exposure during rat brain development. *Alcohol Clin Exp Res.* 1997;21:1418-1428.
23. Gladstone J, Nulman I, Koren G. Reproductive risks of binge drinking during pregnancy. *Reprod Toxicol.* 1996;10:3-13.
24. Floyd RL, Decoufle P, Hungerford DW. Alcohol use prior to pregnancy recognition. *Am J Prev Med.* 1999;17:101-107.
25. Morse J. Women on a binge. *Time.* April 1, 2002; 159(13):56-61.
26. Centers for Disease Control and Prevention. Alcohol use among women of childbearing age—United States, 1991-1999. *MMWR Morb Mortal Wkly Rep.* 2002;51:273-276.
27. Wechsler H, Lee JE, Kuo M, Seibring M, Nelson TF, Lee H. Trends in college binge drinking during a period of increased prevention efforts. *J Am Coll Health.* 2002;50:203-217.
28. Alcohol Epidemiology Program. *Alcohol Policies in the United States: Highlights From the 50 States*. Minneapolis: University of Minnesota; 2000.
29. American Academy of Pediatrics. Alcohol use and abuse: a pediatric concern. *Pediatrics.* 2001;108:185-189.
30. Midanik LT. Validity of self-reported alcohol use: a literature review. *Br J Addict.* 1988;83:1019-1030.
31. Feunekes G, van't Veer P, van Staveren W, et al. Alcohol intake assessment: the sober facts. *Am J Epidemiol.* 1999;150:105-112.
32. Smith PF, Remington PL, Williamson DF, et al. A comparison of alcohol sales data with survey data on self-reported alcohol use in 21 states. *Am J Public Health.* 1990;80:309-312.
33. Nelson DE. Reliability and validity of measures from the Behavioral Risk Factor Surveillance System (BRFSS). *Soc Prev Med.* 2001;46(suppl 1):S3-S42.
34. Wechsler H, Dowdall GW, Maenner G, Gledhill-Hoyt J, Lee H. Changes in binge drinking and related problems among American college students between 1993 and 1997: results of the Harvard School of Public Health college alcohol survey. *J Am Coll Health.* 1998;47:57-68.
35. Liu S, Siegel PZ, RD B, et al. Prevalence of alcohol-impaired driving. *JAMA.* 1997;277:122-125.
36. Wagenaar AC, Wolfson M. Deterring sales and provision of alcohol to minors: a study of enforcement in 295 counties in four states. *Public Health Rep.* 1995;110:419-427.
37. Stout EM, Sloan FA, Liang L, Davies HH. Reducing harmful alcohol-related behaviors: effective regulatory methods. *J Stud Alcohol.* 2000;61:402-412.
38. McKnight AJ, Streff FM. The effect of enforcement upon service of alcohol to intoxicated patrons of bars and restaurants. *Accid Anal Prev.* 1994;26:79-88.
39. Holder HD, Gruenewald PJ, Ponicki WR, et al. Effect of community-based interventions on high-risk drinking and alcohol-related injuries. *JAMA.* 2000; 284:2341-2347.
40. Center for Disease Control and Prevention. Motor-vehicle occupant injury: strategies for increasing use of child safety seats, increasing use of safety belts, and reducing alcohol impaired driving. *MMWR Morb Mortal Wkly Rep.* 2001;50(RR 07):1-13.
41. Chaloupka FJ, Saffer H, Grossman M. Alcohol-control policies and motor-vehicle fatalities. *J Legal Stud.* 1993;22:161-186.
42. Partnership for Prevention. *Alcohol and Health: When Risky Use Means Costly Problems*. Washington, DC: Partnership for Prevention; 2002.
43. Cook PJ, Moore MJ. The economics of alcohol abuse and alcohol-control policies; price levels, including excise taxes, are effective at controlling alcohol consumption; raising excise taxes would be in the public interest. *Health Aff (Millwood).* 2002;21:120-133.
44. Chaloupka FJ, Grossman M, Saffer H. The effects of price on alcohol consumption. *Alcohol Res Health.* 2002;26:22-34.
45. Saffer H, Grossman M. Beer taxes, the legal drinking age, and youth motor vehicle fatalities. *J Legal Stud.* 1987;16:351-374.
46. Rogers JD, Greenfield TK. Beer drinking accounts for most of the hazardous alcohol consumption reported in the United States. *J Stud Alcohol.* 1999; 60:732-739.
47. US Preventive Services Task Force, Screening for Problem Drinking. *Guide to Clinical Preventive Services*. Washington, DC: US Dept of Health and Human Services; 1996:567-582.
48. Coffield AB, Maciosek MV, McGinnis MJ, et al. Priorities among recommended clinical preventive services. *Am J Prev Med.* 2001;21:1-9.
49. Fleming MF, Barry KL, Manwell LB, Johnson K, London R. Brief physician advice for problem drinkers. *JAMA.* 1997;277:1039-1045.
50. Federal Trade Commission. *Self-regulation in the alcohol industry*. Available at: <http://www.ftc.gov/reports/alcohol.alcoholreport.htm>. Accessed October 3, 2002.
51. The Center on Alcohol Marketing and Youth. *OverExposed: youth at target of alcohol advertising in magazines*. Available at <http://camy.org/research/files/overexposed0902.pdf>. Accessed December 10, 2002.
52. Lender ME, Martin JK. *Drinking in America*. New York, NY: The Free Press; 1987.
53. Rae S, ed. *Drink, Drinkers, and Drinking*. London, England: Faber; 1991.