

Pancreatitis

The **pancreas**, an organ located behind the stomach in the upper abdomen, produces enzymes and hormones. The enzymes are discharged through the **pancreatic duct** into the intestine, where they help digest food that has passed through the stomach. The hormones produced by the pancreas, including **insulin** and **glucagon**, are released into the blood and help regulate blood sugar levels. **Pancreatitis** is an inflammation of the pancreas. Pancreatitis is a severe illness with several forms. The June 16, 2004, issue of *JAMA* includes an article about pancreatitis.

RISK FACTORS FOR PANCREATITIS

- Alcohol abuse
- Gallstones
- Abdominal trauma
- Major surgery
- Inherited diseases affecting the pancreas, including **cystic fibrosis**
- High blood levels of calcium (usually due to other medical problems)
- High blood levels of **triglycerides** (a particular type of fat in the blood)

ACUTE PANCREATITIS

Acute pancreatitis typically causes severe abdominal pain, often associated with nausea and vomiting. Because pancreatitis can be severe and life-threatening, antibiotics and hospital care may be required. In severe cases of acute pancreatitis, shock may occur, requiring intensive care and supportive therapy.

GALLSTONE PANCREATITIS

Gallstone pancreatitis is caused by irritation of the pancreas by gallstones in the pancreatic duct. **Jaundice** (yellowing of the skin) may occur with this type of pancreatitis. **Cholecystectomy** (surgical removal of the gallbladder) is necessary to prevent gallstone pancreatitis from occurring again. Because acute gallstone pancreatitis may be a severe illness, several days of supportive care (which may include antibiotics and intravenous fluids in the hospital) may be required before a person is ready to have the gallbladder removed.

CHRONIC PANCREATITIS


Chronic pancreatitis is inflammation and scarring of the pancreas that occurs over a long period. Heavy alcohol use is a common cause. Persons with chronic pancreatitis may have abdominal pain that is either constant or **episodic** (comes and goes). Scarring of the pancreas may lead to failure to produce enough enzymes to digest food and a lack of insulin leading to diabetes. Persons who have acute or chronic pancreatitis may develop a complication called a **pseudocyst**. This is a fluid collection in or around the pancreas. It may push on or cause pressure on other abdominal organs, creating other symptoms. These pseudocysts can also become infected, causing serious illness.

Janet M. Torpy, MD, Writer

Cassio Lynn, MA, Illustrator

Richard M. Glass, MD, Editor

The JAMA Patient Page is a public service of *JAMA*. The information and recommendations appearing on this page are appropriate in most instances, but they are not a substitute for medical diagnosis. For specific information concerning your personal medical condition, *JAMA* suggests that you consult your physician. This page may be photocopied noncommercially by physicians and other health care professionals to share with patients. Any other print or online reproduction is subject to AMA approval. To purchase bulk reprints, call 718/946-7424.



FOR MORE INFORMATION

- American Gastroenterological Association
301/654-2055
www.gastro.org
- National Institute of Diabetes and Digestive and Kidney Diseases
www.niddk.nih.gov
- National Pancreas Foundation
866/726-2737
www.pancreasfoundation.org

INFORM YOURSELF

To find this and previous JAMA Patient Pages, go to the Patient Page link on *JAMA*'s Web site at www.jama.com. Many are available in English and Spanish.

Sources: American Gastroenterological Association, National Institute of Diabetes and Digestive and Kidney Diseases, National Pancreas Foundation

